

**PHIL 1321**

**University Of Houston  
Garson**

510-7F-0007

*ALL COPYRIGHTED MATERIAL HAS BEEN RE-PRINTED BY PERMISSION  
OF THE RIGHTS HOLDER. FOR ONE TIME USE ONLY.*

**Copyright © 2009 University of Houston. All Rights Reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanism, photocopying, recording, or otherwise without prior written permission of the University of Houston c/o University Copy Center, 4810 Calhoun, Ste. 102 , Houston, TX 77004**

**Logic I Study Guide (to help study for quizzes and final)**  
**(The website on WebCT has a more complete study guide.)**

**Quiz 1 Reading: Ch. 1-2 of Propositional Logic**

**Exercises:** Ch. 1: 1, 2 Ch. 2: 1 2 3a, b, c, 5 7 8 9 11

**Concepts:** Argument, Premise, Conclusion, Statement, Validity, Soundness, Inductive Logic, Deductive Logic, Modus Ponens, Conditional, Antecedent, Consequent

**Skills:**

- \* Distinguish validity of an argument from truth of a statement.
- \* Give the two definitions of validity.
- \* Recognize the conclusion and premises of arguments in English using indicator words, on p. 17, Propositional Logic, and their synonyms.
- \* Complete simple proofs using  $\rightarrow$ Out.

**Quiz 2 Reading: Ch. 3-4 of Propositional Logic**

**Exercises:** Ch. 3: 1a, b, c, f, g, j 2b, d 3b, d 6 7 8 9 12 13

Ch. 4: 1b, c, e, f, g, i, j, n 4a, b, c, d 6 9a 10 13 14 16

**Concepts:** Ampersand, Conjunction, Conjunct, Subproof

**Skills:**

- \* Recognize main connectives of complex formulas.
- \* Complete proofs using  $\rightarrow$ ,  $\&$ , rules.
- \* Employ correct strategy for creating subproofs for the  $\rightarrow$ In rule.
- \* Calculate dependency lists.

**Quiz 3 Reading: Ch. 5-7 of Propositional Logic**

**Exercises:** Ch. 5: 1a, b, c, e, f, g, j, m 3a, b, c 4 5 8 9 10 12 13

Ch. 6: 1b, d, f, g 2b, c 5 7 9 10 11 13 15

Ch. 7: 1a, b, e, f, j, l, n, p, r 3a, b 4 5 6 8 10 11 13 14

**Concepts:** Indirect Proof, Reductio ad Absurdum, Standard Contradiction, Disjunction, Disjunct, Exclusive or, Inclusive or, Dilemma

**Skills:**

- \* Accurately diagnose the scope of negation in translating English statements
- \* Construct proofs using the negation rules (-In and -Out)
- \* Understand and appropriately apply the strategies for the  $\leftrightarrow$  Rules.
- \* Understand and apply the strategies for the  $\vee$  Rules, especially the  $\vee$ Out strategy. That is, when  $A \vee B$  appears in the proof, and your last goal is G, know to set the "missing conditionals"  $A \rightarrow G$  and  $B \rightarrow G$  as new goals, and derive G by  $\vee$ Out.

**Quiz 4 Reading: Ch. 8.1, 8.2, 9.2 of Propositional Logic**

**Exercises:** Ch. 8: 2b 4 6 8 9 10 12 13

Ch. 9: 8a, b 9b, f, g 10b, d 11 12 14b, d 15 16 21b, c, i

**Concepts:** Derived Rule, Consistency of Proof Rules, Completeness of Proof Rules

**Skills:**

- \* Know how to demonstrate that a rule is derivable.
- \* Know the Highly Recommended derived rules.
- \* Appropriately employ the strategies for using the Highly Recommended rules that is, use AR and DM to simplify sentences with the forms:  $\neg(A \vee B)$ ,  $\neg(A \rightarrow B)$ ,  $\neg(A \& B)$ ,  $A \vee B$ .
- \* Know how to handle a disjunction as a goal by either converting to  $\rightarrow$  or using the DeMorgan strategy.
- \* Be able to complete more difficult proofs.

**Quiz 5 Reading: Ch. 10, Appendix 4, Propositional Logic,**

"How to Make a Tree", Notes pp. 17-24

**Exercises: Do with Truth Tables: Ch. 10: 1a 4 5 6 8**

**Do with Single Sided trees: Notes: Problems 1-5 and 1-4 at end of "How to Make a Tree" p. 25, 26, Ch. 9: 3 12 Ch. 10: 1c 2-6, 8 10 17 19a 21 24**

**Concepts: Counterexample**

**Skills:**

- \* Calculate the truth table for a statement.
- \* Calculate the truth table for an argument and determine its validity status.
- \* Know how to apply all the tree rules.
- \* Construct trees for arguments and diagnose their validity status.

**Quiz 6 Reading: Ch. 12-13 of Propositional Logic and Ch 2 of Predicate Logic**

**Exercises: Ch. 12: 1 2 4 5 7 9 10**

**Ch. 13: 2 3 4b 6 7 9**

**Predicate Logic, Ch. 2: 1 2 3 4 6 (all parts)**

**Concepts: Valid (TP), Contradiction (TP), Contingent (T), Logical Truth (TP), Tautology (TP), Equivalence (TP), Entailment (TP), Consistency (T).**

**Skills:**

- \* Know how to test all concepts listed above which are marked with "T" with truth tables and trees. Be able to test all concepts marked "P" with proofs.
- \* Be able to distinguish singular from general terms. Especially, be able to diagnose such sentences as "A dog got into the garbage" as general, and "The dog got into the garbage" as singular.
- \* Be able to translate simple predicate logic sentences with the forms Some A are B; All A are B; No A are B; Not all A are B.

**Quiz 7 Reading: Ch 3-4 of Predicate Logic**

**Exercises: Predicate Logic Ch. 3: 1-6, 8-13 do proofs and trees**

**Predicate Logic Ch. 4: 10-15 do proofs and trees**

**Concepts: Instance**

**Skills:**

- \* Complete proofs and trees using the predicate logic rules  $\forall O$ ,  $\exists O$  and  $QE$ .
- \* Order the use of the predicate logic rules properly:  $QE$ ,  $\exists O$ ,  $\forall O$ .
- \* Understand, and avoid breaking, the restriction on the  $\exists O$  rule.

**Final : you should know all of the above plus:**

**Concepts**

Explain the relationship between validity and the following notions:

- \* Counterexample (A valid argument has no counterexample),
- \* Consistency (An argument is invalid if its premises and negated conclusion are consistent)
- \* Closed Tree (An argument is valid if the tree formed from the premises and the negated conclusion is closed in every branch).

**Skills**

- \* Remember all expressions on the Translation Charts in Notes p. 4 and 32.
- \* Recognize main connectives, and apply rules only to main connectives.
- \* Remember perfectly the 10 primitive rules on p. 272, Propositional Logic, the rules  $DM$ ,  $AR$ , and  $DN$ , and the quantifier rules  $QE$ ,  $\exists O$  and  $\forall O$ .
- \* Use the backwards proof finding strategy. Make provisional assumptions (PA) only when appropriate, i.e. only when setting up  $\rightarrow IN$ ,  $-IN$  or  $-OUT$ .
- \* Know, and be able to use, the strategy suggestions on p. 115, Propositional Logic.
- \* Remember truth tables perfectly, and be able to calculate truth tables quickly.
- \* Know the tree rules (Propositional Logic, p. 233-234) and be able to construct trees and counterexamples.
- \* Be able to construct a tree in predicate logic quickly.

# Exercise Answers CH-2

1.1

1. T Snow is white and grass is green

T Snow is white

F Snow is white and grass is black

T Snow is white

F Snow is purple and grass is black

F Snow is purple

you can't get an argument with this form that has a true premise and a false conclusion because the argument form is valid.

2. T Snow is white and grass is green

T Snow is white and polar bears are white

F Snow is white and grass is black

T Snow is white and polar bears are white

F Snow is white and grass is black

F Snow is white and polar bears are purple

T Snow is white and grass is green

F Snow is white and polar bears are purple

## Ch.2

- 1 a.  $I \rightarrow P$  b.  $O \rightarrow W$  c.  $L \rightarrow C$  d.  $S \rightarrow A$  e.  $W \rightarrow P$  f.  $Q \rightarrow P$   
 g.  $B \rightarrow C$  h.  $R \rightarrow B$  i.  $F \rightarrow W$  j.  $A \rightarrow (D \rightarrow T)$  k.  $Q \rightarrow (C \rightarrow R)$  l.  $F \rightarrow (R \rightarrow L)$

2 c. If salt's being added results in the solution's boiling point dropping then if salt is added to the solution then the solution boils sooner

- 3 a. (4) B 1, 3  $\rightarrow$  D (5) C  $\rightarrow$  D 2, 4  $\rightarrow$  D  
 3 c. (4) I  $\rightarrow$  J 1, 2  $\rightarrow$  D (5) J 4, 3  $\rightarrow$  D

7. (1)  $A \rightarrow (A \rightarrow B)$  A  
 (2) A A  
 (3)  $A \rightarrow B$  1, 2  $\rightarrow$  D  
 (4) B 3, 2  $\rightarrow$  D

8. (1) H A  
 (2)  $H \rightarrow R$  A  
 (3)  $R \rightarrow C$  A  
 (4)  $C \rightarrow F$  A  
 (5) R 2, 1  $\rightarrow$  D  
 (6) C 3, 5  $\rightarrow$  D  
 (7) F 4, 6  $\rightarrow$  D

11. (1)  $(F \rightarrow G) \rightarrow (G \rightarrow H)$  A  
 (2)  $F \rightarrow G$  A  
 (3) F A  
 (4)  $G \rightarrow H$  1, 2  $\rightarrow$  D  
 (5) G 2, 3  $\rightarrow$  D  
 (6) H 4, 5  $\rightarrow$  D

## Propositional Logic Translation Chart

**A → B**

If A, (then) B

B if A

provided that  $A, B$

B provided that A

A only if B

A is sufficient for B

B is necessary for A

**A & B**

(Both) A and B

A but B

A however B

A although B

A nevertheless B

A moreover B

A yet B

A even though B

### Antecedent Indicators:

if \_

provided that \_

\_ is sufficient

should \_

### Consequent Indicators:

then \_

only if \_

\_ is necessary

results in \_

brings about \_

leads to \_

### Premise Indicators:

Because

Since

For

### Conclusion Indicators:

Therefore

So

Hence

Thus

Consequently

It follows that

**- A**

Not A

It is false that A

A is mistaken

A is not the case

Negative Prefixes:

Un (unintelligent)

In (indescribable)

Im (immoral)

**A ↔ B**

A if and only if B (A iff B)

A just in case B

A exactly when B

A exactly if B

A but only if B

A is a necessary and sufficient condition for B

**A ∨ B**

(Either) A or B


A unless B

**-(A ∨ B) or -A & -B**


Neither A nor B

# Rules for Boxes (Optional)


1. Every box is headed by 'PA' and 'PA' is used only at the head of a box.  
EXAMPLES:




WRONG!!  
 ('PA' goes at the top of each box)




WRONG!!  
 ('PA' appears without a box in the second line)




2. →In, ~In and -Out close a box, and no other rules do.  
EXAMPLES:



B 2, 1 →out  
WRONG!!  
 (→out does not close a box)




WRONG!!  
 (→In requires that you close the box)



(A & B) → B →IN 1-2  
OK

3. When a box has been closed, no line inside the box can be used.

EXAMPLE:



WRONG!! (Lines 2, 3, and 4 can not be used because the box 2-4 is closed by the time you get to line 6.)

# Strategies for Proof Finding

If you see this:

Then Do This:

$A \rightarrow B$  GOAL

$A \supset PA$   
 $B$

$A \rightarrow B$  GOAL  $\rightarrow$  IN

$A \& B$  GOAL

$A$  GOAL  
 $B$  GOAL  
 $A \& B$  GOAL & IN

$A \leftrightarrow B$  GOAL

$A \rightarrow B$  GOAL  
 $B \rightarrow A$  GOAL  
 $A \leftrightarrow B$  GOAL  $\leftrightarrow$  IN

$A \vee B$  GOAL

Do work elsewhere  
 If nothing else  
 can be done  
 try these...

$A$  GOAL  
 $A \vee B$  GOAL  $\vee$  IN     OR      $B$  GOAL  
 $A \vee B$  GOAL  $\vee$  IN     OR      $\sim A \rightarrow B$  GOAL  
 $A \vee B$  GOAL  $\vee$  IN     OR      $A \vee B$  GOAL AR

$A \rightarrow G$  (AVAILABLE)

$A \rightarrow G$

$\vdots$

$G$  GOAL

$A$  GOAL  
 $G$  GOAL  $\rightarrow$  Out

$A \vee B$  (AVAILABLE)

$A \vee B$

$\vdots$

$G$  GOAL

$A \rightarrow G$  GOAL     OR      $A \vee B$  (PROVEN)  
 $B \rightarrow G$  GOAL     OR      $\sim A \rightarrow B$  (Arrow) AR  
 $G$  GOAL  $\vee$  OUT     OR      $G$  GOAL

$\sim (A \& B)$      convert to  $A \rightarrow \sim B$      (Arrow)     AR

$\sim (A \rightarrow B)$      convert to  $A \& \sim B$      (Arrow)     AR

$\sim (A \vee B)$      convert to  $\sim A \& \sim B$      (DeMorgan)     DM

$G$  GOAL  
 (nothing works)

$\sim G \supset PA$   
 $P \& \sim P$

$G$  GOAL I.P. (out)

Try Indirect Proof  
 ?&? stands for  
 a contradiction

## How to choose a Contradiction

$P \& \sim P$  GOAL

1. Prefer long negated formulas already available,
2. Avoid formulas already used as contradictions for I.P.
3. Avoid using GOAL for I.P. as part of the contradiction



## PROOF STRATEGY

0. Can I apply DN, &Out,  $\rightarrow$ Out or  $\leftrightarrow$ Out to lines already available?  
Yes? Then do it.

1. What is my goal?

2. Is my goal  $\mathcal{A} \rightarrow \mathcal{B}$ ,  $\mathcal{A} \& \mathcal{B}$ , or  $\mathcal{A} \leftrightarrow \mathcal{B}$ ?

Yes? Then assume it will come by the corresponding In rule. For example, if it is  $\mathcal{A} \rightarrow \mathcal{B}$ , make a box with  $\mathcal{A}$  a PA, and set  $\mathcal{B}$  as a new goal.

If it is  $\mathcal{A} \& \mathcal{B}$ , set  $\mathcal{A}$  and  $\mathcal{B}$  as new goals.

If it is  $\mathcal{A} \leftrightarrow \mathcal{B}$ , set  $\mathcal{A} \rightarrow \mathcal{B}$  and  $\mathcal{B} \rightarrow \mathcal{A}$  as new goals.

3. Is my goal  $\mathcal{A} \vee \mathcal{B}$ ?

Yes? Then if either  $\mathcal{A}$  or  $\mathcal{B}$  is available, use vIn. Otherwise work elsewhere.

4. Does my goal appear to the right of  $\rightarrow$  in an available line?

Yes? Then set what is to the left of  $\rightarrow$  as a new goal. When you get that goal, use  $\rightarrow$ Out to get what was on the right of  $\rightarrow$ . (Your old goal.)

5. Is  $\mathcal{A} \vee \mathcal{B}$  available?

Yes? Then set up vOut. That means set  $\mathcal{A} \rightarrow G$  and  $\mathcal{B} \rightarrow G$  as new goals, where G is your old goal.

6. Is  $\sim(\mathcal{A} \vee \mathcal{B})$  available?

Yes? Then convert to  $\sim \mathcal{A} \& \sim \mathcal{B}$  with DM.

7. Is  $\sim(\mathcal{A} \rightarrow \mathcal{B})$  or  $\sim(\mathcal{A} \& \mathcal{B})$  available?

Yes? Then use AR to convert this to  $\mathcal{A} \& \sim \mathcal{B}$  or  $\mathcal{A} \rightarrow \sim \mathcal{B}$  respectively.

8. Are you stuck?

Yes? Then use IP. Make a box with the opposite of your goal a PA and chose a contradiction as a new goal. In choosing a contradiction look for negative sentences in lines available. Long negations are a good choice. Do not use the goal for IP as part of your contradiction, and do not use contradictions previously used for IP.

# Exercise Answers Ch 3-4

Ch. 3

1 a. W & S c. S & B g. A & B → F

2 d If Miami wins its last game then both N.Y loses its last game and Miami wins the championship.

- 3 d (1) (H & I) → (J & K) A  
 (2) I & H A  
 (3) I 2 & 0  
 (4) H 2 & 0  
 (5) H & I 4, 3 & I  
 (6) J & K 1, 5 → 0  
 (7) J 6 & 0

6. (1) K A  
 (2) S A  
 (3) (K & S) → H A  
 (4) K & S 1, 2 & I  
 (5) H 3, 4 → 0

7. (1) I & F A  
 (2) I → (F → C) A  
 (3) I 1 & 0  
 (4) F → C 2, 3 → 0  
 (5) F 1 & 0  
 (6) C 4, 5 → 0

12. (1) E A  
 (2) M & V A  
 (3) E → (V & M) → H A  
 (4) H → I A  
 (5) (V & M) → H 3, 1 → 0  
 (6) M 2 & 0  
 (7) V 2 & 0  
 (8) (V & M) 7, 6 & I n  
 (9) H 5, 8 → 0  
 (10) I 4, 9 → 0

9. (1) (F & R) & P A  
 (2) F & R 1 & 0  
 (3) F 2 & 0  
 (4) R 2 & 0  
 (5) P 1 & 0  
 (6) R & P 4, 5 & I  
 (7) F & (R & P) 3, 6 & I

Ch. 4

1. c. P → A e. S → R g. R → D i. (W & D) → (M → R)

- 4 a. (3) B 1, 2 → 0  
 (4) A & B 2, 3 & I

- 4 b. (2) G & F (PA)  
 (3) F 2 & 0  
 (4) G → H 1, 3 → 0  
 (5) G 2 & 0  
 (6) H 4, 5 → 0  
 (7) (G & F) → H 2-6 → I

- 4 d. (2) F (PA)  
 (3) G (PA)  
 (4) G & F 2, 3 & I n  
 (5) H 1, 4 → 0  
 (6) G → H 3-5 → I  
 (7) F → (G → H) 2-6 → I n

- 9 a. (1) S → P A  
 (2) (P → N) & (N → C) A  
 (3) S (PA)  
 (4) P → N 2 & 0  
 (5) N → C 2 & 0  
 (6) P 1, 3 → 0  
 (7) N 4, 6 → 0  
 (8) C 5, 7 → 0  
 (9) S → C 3-8 → I

13. (1) (T → L) & (T → M) A  
 (2) T (PA)  
 (3) T → L 1 & 0  
 (4) T → M 1 & 0  
 (5) L 3, 2 → 0  
 (6) M 4, 2 → 0  
 (7) L & M 5, 6 & I  
 (8) T → (L & M)

14. (1) T → (L & M)  
 (2) T (PA)  
 (3) L & M 1, 2 → 0  
 (4) L 3 & 0  
 (5) T → L 2-4 → I  
 (6) T (PA)  
 (7) L & M 1, 6 → 0  
 (8) M 7 & 0  
 (9) T → M 6-8 → I

16. (1) D → S  
 (2) S → T  
 (3) (T & S) → C  
 (4) ((S & T) & C) → R  
 (5) D (PA)  
 (6) S 1, 5 → 0  
 (7) T 2, 6 → 0  
 (8) T & S 7, 6 & I  
 (9) C 3, 8 → 0  
 (10) S & T 6, 7 & I  
 (11) (S & T) & C 10, 9 & I  
 (12) R 4, 11 → 0  
 (13) D → R 5-12 → I

Ignore the answer for 14 given in the book. It is not compatible with the box method.

# Exercise Answers Ch 5

Ch. 5

1 a.  $\neg T$  1c.  $\neg F \rightarrow A$  1e.  $\neg P \& W$  1g.  $(\neg C \& \neg W) \& \neg M$   
 or  $\neg C \& (\neg W \& \neg M)$

1m.  $(\neg T \& A) \rightarrow (\neg H \& \neg P)$

3a. 1 (1)  $\neg A \rightarrow B$  A  
 2 (2)  $\neg A \rightarrow \neg B$  A  
 3 (3)  $\neg A$  (PA)  
 1,3 (4) B 1,3  $\rightarrow$  0  
 2,3 (5)  $\neg B$  2,3  $\rightarrow$  0  
 1,2,3 (6) B &  $\neg B$  4,5 & I  
 1,2 (7) A 3-6 -O

3c. 1 (1)  $E \rightarrow F$  A  
 2 (2)  $\neg(E \& F)$  A  
 3 (3) E (PA)  
 1,3 (4) F 1,3  $\rightarrow$  0  
 1,3 (5)  $E \& F$  3,4 & I  
 1,2,3 (6)  $E \& F \& \neg(E \& F)$  5,2 & I  
 1,2 (7)  $\neg E$  3-6 -I

4. 1 (1)  $P \rightarrow M$  A  
 2 (2)  $\neg M$  A  
 3 (3) P (PA)  
 1,3 (4) M 1,3  $\rightarrow$  0  
 1,2,3 (5)  $M \& \neg M$  4,2 & I  
 1,2 (6)  $\neg P$  3-5 -I

8. 1 (1) R  
 2 (2) L  
 3 (3)  $(S \& R) \rightarrow \neg L$   
 4 (4) S (PA)  
 1,4 (5)  $S \& R$  4,1 & I  
 1,3,4 (6)  $\neg L$  3,5  $\rightarrow$  0  
 1,2,3,4 (7)  $L \& \neg L$  3,6 & I  
 1,2,3 (8)  $\neg S$  4-7 -I

10. 1 (1)  $A \rightarrow S$  A  
 2 (2)  $G \rightarrow A$  A  
 3 (3)  $S \rightarrow \neg G$  A  
 4 (4) G (PA)  
 2,4 (5) A 2,4  $\rightarrow$  0  
 1,2,4 (6) S 1,5  $\rightarrow$  0  
 1,2,3,4 (7)  $\neg G$  3,6  $\rightarrow$  0  
 1,2,3,4 (8)  $G \& \neg G$  4,7 & I  
 1,2,3 (9)  $\neg G$  4-8 -I

12. 1 (1)  $L \rightarrow T$  A  
 2 (2) J A  
 3 (3)  $\neg(T \& J)$  A  
 4 (4) L (PA)  
 1,4 (5) T 1,4  $\rightarrow$  0  
 1,2,4 (6)  $T \& J$  5,2 & I  
 1,2,3,4 (7)  $(T \& J) \& \neg(T \& J)$  3,6 & I  
 1,2,3 (8)  $\neg L$  4-7 -I

# Exercise Answers Ch 6

Ch 6 1d.  $\neg A \leftrightarrow R$  1g.  $D \leftrightarrow H$  2c. Both Smith wins the batting crown if and only if Smith gets a hit, and Jones makes an out.

7. (1)  $(C \rightarrow L) \& (L \rightarrow F)$  A  
 2 (2)  $F \rightarrow C$  A  
 3 (3) 

C	(PA)
$C \rightarrow L$	1 & 0
L	4, 3 $\rightarrow$ 0
$L \rightarrow F$	1 & 0
F	6, 5 $\rightarrow$ 0

  
 1 (4) 

L	4, 3 $\rightarrow$ 0
$L \rightarrow F$	1 & 0

  
 1, 3 (5) 

F	6, 5 $\rightarrow$ 0
---	----------------------

  
 1 (6)  $C \rightarrow F$  3-7  $\rightarrow$  I  
 1, 2 (9)  $C \leftrightarrow F$  8, 2  $\leftrightarrow$  I

10. (1)  $I \rightarrow (B \leftrightarrow M)$  A  
 2 (2) B A  
 3 (3)  $M \rightarrow T$  A  
 4 (4)  $\neg T$  A  
 5 (5) 

I	(PA)
$B \leftrightarrow M$	4, 5 $\rightarrow$ 0
$B \rightarrow M$	6 $\leftrightarrow$ 0
M	7, 2 $\rightarrow$ 0
T	3, 8 $\rightarrow$ 0
$T \& \neg T$	9, 4 & I

  
 1, 5 (6)  $B \leftrightarrow M$  4, 5  $\rightarrow$  0  
 1, 5 (7)  $B \rightarrow M$  6  $\leftrightarrow$  0  
 1, 2, 5 (8) M 7, 2  $\rightarrow$  0  
 1, 2, 3, 5 (9) T 3, 8  $\rightarrow$  0  
 1, 2, 3, 4, 5 (10)  $T \& \neg T$  9, 4 & I  
 1, 2, 3, 4 (11)  $\neg I$  5-10 -In

11. (1)  $N \rightarrow R$  A  
 2 (2)  $O \leftrightarrow R$  A  
 3 (3)  $(O \rightarrow R) \rightarrow L$  A  
 2 (4)  $O \rightarrow R$  2  $\leftrightarrow$  0  
 2, 3 (5) L 3, 4  $\rightarrow$  0  
 6 (6) 

N	(PA)
R	1, 6 $\rightarrow$ 0
$R \rightarrow O$	2 $\leftrightarrow$ 0
O	8, 7 $\rightarrow$ 0

  
 1, 6 (7) 

R	1, 6 $\rightarrow$ 0
$R \rightarrow O$	2 $\leftrightarrow$ 0

  
 2 (8) 

O	8, 7 $\rightarrow$ 0
---	----------------------

  
 1, 2, 6 (9)  $N \rightarrow O$  6-9  $\rightarrow$  I  
 1, 2 (10)  $(N \rightarrow O) \& L$  10, 5 & I  
 1, 2, 3 (11)  $(N \rightarrow O) \& L$  10, 5 & I

15. (1)  $A \leftrightarrow B$  A  
 2 (2)  $\neg(A \& B)$  A  
 3 (3) 

A	(PA)
$A \rightarrow B$	1 $\leftrightarrow$ 0
B	4, 3 $\rightarrow$ 0
$A \& B$	3, 5 & In
$(A \& B) \& \neg(A \& B)$	2, 6 & I

  
 1 (4) 

A	(PA)
$A \rightarrow B$	1 $\leftrightarrow$ 0
B	4, 3 $\rightarrow$ 0
$A \& B$	3, 5 & In
$(A \& B) \& \neg(A \& B)$	2, 6 & I

  
 1, 3 (5) 

A	(PA)
$A \rightarrow B$	1 $\leftrightarrow$ 0
B	4, 3 $\rightarrow$ 0
$A \& B$	3, 5 & In
$(A \& B) \& \neg(A \& B)$	2, 6 & I

  
 1, 3 (6) 

A	(PA)
$A \rightarrow B$	1 $\leftrightarrow$ 0
B	4, 3 $\rightarrow$ 0
$A \& B$	3, 5 & In
$(A \& B) \& \neg(A \& B)$	2, 6 & I

  
 1, 2, 3 (7) 

A	(PA)
$A \rightarrow B$	1 $\leftrightarrow$ 0
B	4, 3 $\rightarrow$ 0
$A \& B$	3, 5 & In
$(A \& B) \& \neg(A \& B)$	2, 6 & I

  
 1, 2 (8)  $\neg A$  3-7 -In  
 9 (9) 

B	(PA)
$B \rightarrow A$	1 $\leftrightarrow$ 0
A	10, 9 $\rightarrow$ 0
$A \& B$	11, 9 & I
$(A \& B) \& \neg(A \& B)$	12, 2 & I

  
 1 (10) 

B	(PA)
$B \rightarrow A$	1 $\leftrightarrow$ 0
A	10, 9 $\rightarrow$ 0
$A \& B$	11, 9 & I
$(A \& B) \& \neg(A \& B)$	12, 2 & I

  
 1, 9 (11) 

B	(PA)
$B \rightarrow A$	1 $\leftrightarrow$ 0
A	10, 9 $\rightarrow$ 0
$A \& B$	11, 9 & I
$(A \& B) \& \neg(A \& B)$	12, 2 & I

  
 1, 9 (12) 

B	(PA)
$B \rightarrow A$	1 $\leftrightarrow$ 0
A	10, 9 $\rightarrow$ 0
$A \& B$	11, 9 & I
$(A \& B) \& \neg(A \& B)$	12, 2 & I

  
 1, 3, 9 (13) 

B	(PA)
$B \rightarrow A$	1 $\leftrightarrow$ 0
A	10, 9 $\rightarrow$ 0
$A \& B$	11, 9 & I
$(A \& B) \& \neg(A \& B)$	12, 2 & I

  
 1, 2 (14)  $\neg B$  9-13 -In  
 1, 2 (15)  $\neg A \& \neg B$  8, 14 & I

# Exercise Answers Ch 7

Ch 7

1a. MVD 1e.  $(S \rightarrow B) \vee L$  1l.  $(R \vee A) \& \sim (R \& A)$

1p.  $((B \& G) \vee (\sim B \& \sim G)) \rightarrow P$

4. Line is obtained from 5. But 5 is in a closed box so the step is illegal.

3a. 1 (1)  $\sim (A \vee B)$  A  
 2 (2)  $A \& B$  (PA)  
 2 (3) A 2 & O  
 2 (4)  $A \vee B$  3 v In  
 1, 2 (5)  $(A \vee B) \& \sim (A \vee B)$  4, 1 & In  
 1 (6)  $\sim (A \& B)$  2-5 -In

5. (1)  $B \vee D$  A  
 (2)  $B \rightarrow F$  A  
 (3)  $O \rightarrow F$  A  
 (4) F 1, 2, 3 v O.

6. (1)  $A \& B$  A  
 (2) A 1 & O  
 (3)  $C \vee A$  2 v I  
 (4)  $(C \vee A) \vee D$  3 v I

8. 1 (1) C A  
 2 (2)  $C \rightarrow (J \vee A)$  A  
 3 (3)  $J \rightarrow \sim S$  A  
 4 (4)  $A \rightarrow \sim S$  A  
 5 (5)  $\sim M \rightarrow S$  A

11. 1 (1)  $L \vee S$  A  
 2 (2)  $(L \rightarrow K) \& (S \rightarrow C)$  A  
 3 (3)  $(K \rightarrow G) \& (C \rightarrow G)$  A  
 2 (4)  $L \rightarrow K$  2 & O  
 2 (5)  $\sim S \rightarrow C$  2 & O  
 3 (6)  $K \rightarrow G$  3 & O  
 3 (7)  $C \rightarrow G$  3 & O

6 (6)  $\sim M$  (PA)  
 5, 6 (7) S 5, 6  $\rightarrow$  O  
 1, 2 (8)  $J \vee A$  2, 1  $\rightarrow$  O  
 1, 2, 3, 4 (9)  $\sim S$  8, 9, 4 v O  
 1, 2, 3, 4, 5, 6 (10)  $S \& \sim S$  7, 9 & I  
 1, 2, 3, 4, 5 (11) M - O 6 - 10

8 (8) L (PA)  
 2, 8 (9) K 4, 8  $\rightarrow$  O  
 2, 3, 8 (10) G 6, 9  $\rightarrow$  O

13. 1 (1)  $(A \& B) \vee (A \& C)$  A

2, 3 (11)  $L \rightarrow G$  8-10  $\rightarrow$  I

2 (2) A & B (PA)  
 2 (3) A 2 & O  
 2 (4) B 2 & O  
 2 (5)  $B \vee C$  4 v In  
 2 (6)  $A \& (B \vee C)$  3, 5 & In

12 (12) S (PA)  
 2, 12 (13) C 5, 12  $\rightarrow$  O  
 2, 3, 12 (14) G 7, 13  $\rightarrow$  O

(7)  $(A \& B) \rightarrow (A \& (B \vee C))$  2-6  $\rightarrow$  I

2, 3 (15)  $S \rightarrow G$  12-14  $\rightarrow$  I

8 (8) A & C (PA)  
 8 (9) A 8 & O  
 8 (10) C 8 & O  
 8 (11)  $B \vee C$  10 v I  
 8 (12)  $A \& (B \vee C)$  9, 11 & I

1, 2, 3 (16) G 1, 11, 15 v O

(13)  $(A \& C) \rightarrow (A \& (B \vee C))$  8-12  $\rightarrow$  I

1 (14)  $A \& (B \vee C)$  1, 7, 13 v O

Exercise Answers Ch 8

Ch 8 26 IGNORE THE ANSWER IN THE BOOK

2b 1 (1) TVP A

2 (2) ~P A

3 (3) T (PA)

(4)  $T \rightarrow T$  3-3  $\rightarrow I$

5 (5) P (PA)

6 (6) ~T (PA)  
2,6 (7) P & ~P 2,5 & In

2 (8) T -0 6-8

2 (9)  $P \rightarrow T$  5-8  $\rightarrow I$

1,2 (10) T 1,4,9 VOUT

T	(PA)
T & ~P	2,3 & In
T	3a 20

10, (1) T A

2 (2)  $(T \& 0) \rightarrow M$  A

3 (3) ~M A

4 (4) -0  $\rightarrow$  (C&R) A

5 (5) 0 (PA)

1,5 (6) T & 0 5,1 & I

1,2,5 (7) M 2,6  $\rightarrow$  0

1,2,3,5 (8) M & ~M 7,3 & I

1,2,3 (9) -0 -I 5-8

1,2,3,4 (10) C&R 4,9  $\rightarrow$  0

1,2,3,4 (11) -0 & (C&R) 9,10 & I

6. 1 (1) A  $\rightarrow$  R A

2 (2) R  $\rightarrow$  G A

3 (3) G  $\rightarrow$  (F&E) A

4 (4) A (PA)

1,4 (5) R 1,4  $\rightarrow$  0

1,2,4 (6) G 2,5  $\rightarrow$  0

1,2,3,4 (7) F&E 3,6  $\rightarrow$  0

4,2,3,4 (8) G & (F&E) 6,7 & In

1,2,3,4 (9) R & (G & (F&E)) 5,8 & I

1,2,3 (10) A  $\rightarrow$  (R & (G & (F&E))) 4-9  $\rightarrow$  I

9. 1 (1) (ARP) v (~AR-C) A

2 (2) ARP PA

2 (3) P 2 & 0

2 (4) PV-C 3 VI

(5) (ARP)  $\rightarrow$  (PV-C) 2-4  $\rightarrow$  I

6 (6) ~AR-C (PA)

6 (7) ~C 6 & 0

6 (8) PV-C 7 VI

(9) (~AR-C)  $\rightarrow$  (PV-C) 6-8  $\rightarrow$  I

1 (10) PV-C 1,5,9 V0

13. 1 (1) FVI A

2 (2) F  $\rightarrow$  (~D  $\rightarrow$  N) A

3 (3) ~D A

4 (4) I  $\rightarrow$  P A

5 (5) P  $\rightarrow$  T A

6 (6) T  $\rightarrow$  (~D  $\rightarrow$  (B  $\rightarrow$  N)) A

7 (7) B A

8 (8) F (PA)

2,8 (9) ~D  $\rightarrow$  N 2,8  $\rightarrow$  0

2,3,8 (10) N 9,3  $\rightarrow$  0

2,3 (11) F  $\rightarrow$  N 8  $\rightarrow$  10  $\rightarrow$  I

12 (12) I (PA)

4,12 (13) P 4,12  $\rightarrow$  0

4,5,12 (14) T 5,13  $\rightarrow$  0

4,5,6,12 (15) ~D  $\rightarrow$  (B  $\rightarrow$  N) 6,14  $\rightarrow$  0

3,4,5,6,12 (16) B  $\rightarrow$  N 15,3  $\rightarrow$  0

3,4,5,6,7,12 (17) N 16,7  $\rightarrow$  0

3,4,5,6,17 (18) I  $\rightarrow$  N 12-17  $\rightarrow$  I

(19) N 1,11,18 V0

# A Graphical Summary of the Rules of Propositional Logic

IN RULES

OUT RULES

	IN RULES	OUT RULES
$\rightarrow$	<div style="border: 1px solid black; padding: 5px; display: inline-block;"> <math>a</math>  <math>:</math>  <math>B</math> </div> <div style="text-align: right; margin-top: 5px;"> <span style="border: 1px solid black; border-radius: 50%; padding: 2px;">PA</span> </div> <hr style="width: 80%; margin: 5px auto;"/> $a \rightarrow B \quad \rightarrow \text{IN}$	$a \rightarrow B$ $a$ <hr style="width: 80%; margin: 5px auto;"/> $B \quad \rightarrow \text{OUT}$
$\&$	$a$ $B$ <hr style="width: 80%; margin: 5px auto;"/> $a \& B \quad \& \text{IN}$	$a \& B$ <hr style="width: 80%; margin: 5px auto;"/> $a \quad \& \text{OUT}$ $a \& B$ <hr style="width: 80%; margin: 5px auto;"/> $B \quad \& \text{OUT}$
$\vee$	$a$ <hr style="width: 80%; margin: 5px auto;"/> $a \vee B \quad \vee \text{IN}$ $a$ <hr style="width: 80%; margin: 5px auto;"/> $B \vee a \quad \vee \text{IN}$	$a \vee B$ $a \rightarrow C$ $B \rightarrow C$ <hr style="width: 80%; margin: 5px auto;"/> $C \quad \vee \text{OUT}$
$\leftrightarrow$	$a \rightarrow B$ $B \rightarrow a$ <hr style="width: 80%; margin: 5px auto;"/> $a \leftrightarrow B \quad \leftrightarrow \text{IN}$	$a \leftrightarrow B$ <hr style="width: 80%; margin: 5px auto;"/> $a \rightarrow B \quad \leftrightarrow \text{OUT}$ $a \leftrightarrow B$ <hr style="width: 80%; margin: 5px auto;"/> $B \rightarrow a \quad \leftrightarrow \text{OUT}$
$\sim$	<div style="border: 1px solid black; padding: 5px; display: inline-block;"> <math>a</math>  <math>:</math>  <math>? \&amp; - ?</math> </div> <div style="text-align: right; margin-top: 5px;"> <span style="border: 1px solid black; border-radius: 50%; padding: 2px;">PA</span> </div> <hr style="width: 80%; margin: 5px auto;"/> $\sim a \quad \sim \text{IN}$	<div style="border: 1px solid black; padding: 5px; display: inline-block;"> <math>\sim a</math>  <math>:</math>  <math>? \&amp; \sim ?</math> </div> <div style="text-align: right; margin-top: 5px;"> <span style="border: 1px solid black; border-radius: 50%; padding: 2px;">PA</span> </div> <hr style="width: 80%; margin: 5px auto;"/> $a \quad \sim \text{OUT}$

## Chapter 9.

### Highly Recommended Derived Rules (Use TOP-Down)

$$\frac{\sim(A \rightarrow B)}$$

$$A \& \sim B \text{ AR}$$

$$\frac{\sim(A \& B)}$$

$$A \rightarrow \sim B \text{ AR}$$

$$\frac{\sim(A \vee B)}$$

$$\sim A \& \sim B \text{ DM}$$

$$\frac{\sim \sim A}$$

$$A \text{ DN}$$

### Recommended Derived Rules

$$\frac{\sim A \vee B}$$

$$\sim A \rightarrow B \text{ AR}$$

Simplify the problem by getting rid of  $\vee$ . (Top-Down)

$$\frac{\sim A \rightarrow B \text{ GOAL}}{A \vee B \text{ AR}}$$

In case  $A \vee B$  is a goal convert it to  $\sim A \rightarrow B$  which is likely to be easier to obtain.

$$\frac{\sim A \rightarrow \sim B}{B \rightarrow A \text{ CN}}$$

You might also use other versions of CN. But make sure that after you do CN there is some rule you can apply to the resulting conditional. DO NOT apply CN without a way to use the result.



# Exercise Answers Ch 9

Ch 9

8a. (3) -- B 2 DN  
(4) A 1,3 DA

9. g (3) is not the result of Applying DN to (2) -- (P & R) Is.

10d. (1)  $F \rightarrow (L \& A)$  A  
(2)  $\neg L \vee \neg A$  A  
(3)  $\neg(L \& A)$  2, DM.  
(4)  $\neg(LL \& A) \rightarrow \neg F$  1 CN  
(5)  $\neg F$  4,3  $\rightarrow$  D

11... (1)  $K \& (L \& B) \rightarrow \neg K$  A  
(2) K 1 R0  
(3)  $(A \& B) \rightarrow \neg K$  1 R0  
(4)  $K \rightarrow \neg(A \& B)$  3 CN  
(5)  $\neg(A \& B)$  4,2  $\rightarrow$  D  
(6)  $\neg A \vee \neg B$  5 DM.

15.1 (1)  $(\neg S \rightarrow D) \& (S \rightarrow H)$  A  
2 (2)  $H \rightarrow B$  A

14d.1 (1)  $T \vee F$  A

3	(3)	$\neg(D \vee B)$	PA
3	(4)	$\neg D \& \neg B$	3 DM
3	(5)	$\neg D$	4 R0
3	(6)	$\neg B$	4 R0
2,3	(7)	$\neg B \rightarrow \neg H$	2 CN
2,3	(8)	$\neg H$	7,6 $\rightarrow$ D
2,3	(9)	$S \rightarrow H$	1 R0
1	(10)	$\neg H \rightarrow \neg S$	9 CN
4,2,3	(11)	$\neg S$	10,9 $\rightarrow$ D
1	(12)	$\neg S \rightarrow D$	1 R0
1,2,3	(13)	D	12,11 $\rightarrow$ D
1,2,3	(14)	$D \& \neg D$	13,5 & I
1,2	(15)	DVB	-0 3-14

2 (2)  $T \rightarrow \neg C$  A  
3 (3)  $F \rightarrow L$  A

4	(4)	G	PA
2	(5)	$G \rightarrow \neg T$	2 CN
2,4	(6)	$\neg T$	5,4 $\rightarrow$ D
1	(7)	$\neg T \rightarrow F$	1 AR
4,2,4	(8)	F	7,6 $\rightarrow$ D
4,2,3,4	(9)	L	3,8 $\rightarrow$ D
1,2,3,4	(10)	$G \rightarrow L$	4-9 $\rightarrow$ I
1,2,3	(11)	$\neg G \vee L$	10 AR

21 b.1 (1)  $\neg(A \& B)$  A  
2 (2) A A

21 c.1 (1)  $\neg(A \rightarrow B)$  A

3	(3)	B	PA
2,3	(4)	$A \& B$	2,3 & In
1,2,3	(5)	$A \& B \& \neg(A \& B)$	4,2 & I
1,2	(6)	$\neg B$	3-5 -I

2	(2)	$\neg A$	PA
3	(3)	A	PA
4	(4)	$\neg B$	PA
2,3	(5)	$A \& \neg A$	2,3 & I
2,3	(6)	B	4-5 -D
2	(7)	$A \rightarrow B$	3-6 $\rightarrow$ I
4,2	(8)	$(A \rightarrow B) \& \neg(A \rightarrow B)$	7,1 & I
1	(9)	A	2-8 -D

21 c.1 (1)  $A \& \neg B$   
2 (2)  $A \rightarrow B$  PA  
1 (3) A 1 R0  
1,2 (4) B 2,3  $\rightarrow$  D  
1 (5)  $\neg B$  1 R0  
1,2 (6)  $B \& \neg B$  4,5 & I  
1 (7)  $\neg(A \rightarrow B)$  2-6 -I

10	(10)	B	PA
11	(11)	A	PA
10,11	(12)	$A \& B$	11,10 & I
10,11	(13)	B	12 R0
10	(14)	$A \rightarrow B$	11-13 $\rightarrow$ I
1,10	(15)	$(A \rightarrow B) \& \neg(A \rightarrow B)$	14,1 & I
1	(16)	$\neg B$	10-15 -I
1	(17)	$A \& \neg B$	9,16 & I

# Exercise Answers Ch 10

CH 10

1a.

A	A	$\neg\neg A$
T	T	T
F	F	F
*	*	*

VALID

4.

C	$C \rightarrow \sim F$	$\sim F$	$\neg C$
T	F	F	T
F	T	F	F
T	T	T	T
F	F	T	F
*	*	*	*

INVALID

5.  $P \leftrightarrow W$  |  $\neg(P \leftrightarrow W)$  |  $\neg\neg(P \leftrightarrow W)$

P	W	$P \leftrightarrow W$	$\neg(P \leftrightarrow W)$	$\neg\neg(P \leftrightarrow W)$
T	T	T	F	T
F	F	T	F	T
T	F	F	T	F
F	T	F	T	F
*	*	*	*	*

VALID

6.

E	P	T	$E \rightarrow P$	$(E \leftrightarrow P) \rightarrow T$	$\neg(E \leftrightarrow P)$	$\neg\neg(E \leftrightarrow P)$
T	T	T	T	T	F	T
F	T	T	F	F	T	F
T	F	T	F	F	T	F
F	F	T	T	T	F	T
T	T	F	T	T	F	T
F	T	F	F	F	T	F
T	F	F	F	F	T	F
F	F	F	T	T	F	T
*	*	*	*	*	*	*

INVALID

8.  $A \leftrightarrow C$  |  $A \rightarrow C, C \rightarrow A$  |  $(A \leftrightarrow C) \rightarrow E$  |  $\neg(A \leftrightarrow C) \rightarrow E$

A	C	E	$A \rightarrow C$	$C \rightarrow A$	$(A \leftrightarrow C) \rightarrow E$	$\neg(A \leftrightarrow C) \rightarrow E$
T	T	T	T	T	T	T
F	T	T	F	T	F	T
T	F	T	T	F	F	T
F	F	T	T	T	T	T
T	T	F	T	T	F	F
F	T	F	F	T	F	T
T	F	F	T	F	F	T
F	F	F	T	T	T	T
*	*	*	*	*	*	*

VALID

## How to Make a Tree

There is a major defect with the truth table method for determining whether an argument is valid. When there are many different letters in an argument, the number of rows in the truth table can be very large. Since the number of rows doubles with each added letter (for 2 letters - 4 rows, 3 letters - 8 rows, 4 letters - 16 rows, etc..) the table contains more than a thousand rows for an argument with only 10 letters. The tree method is a variation on the truth table method which can vastly shorten validity calculation. True, a tree may take as long as the corresponding table in the worst case, but the chances are very good that it will yield massive savings in effort.

The main idea behind the tree method is to avoid calculating any parts of the table which do not contribute to the main objective in the validity test, namely to find a row where all the premises are T and the conclusion is F. If there is such a row, the argument is invalid, because validity requires that it is impossible for (all) the premises to be T and the conclusion F. Let us call a row with (all) T premises and a F conclusion a counterexample. The idea behind tables is to examine all possible combinations of truth values to see whether there is a counterexample. If there is, the argument is invalid, and if not, it is valid. The tree method is more economical than tables because it calculates only those portions of the table where a counterexample can be. To help illustrate the point, consider this table for the following invalid argument:  $Q \rightarrow \neg P \vdash P \rightarrow Q$ .

P	Q	$Q \rightarrow \neg P$	$\vdash P \rightarrow Q$	
T	T	F	T	
F	T	T	T	
T	F	T	F	← Counterexample
F	F	T	T	

Note that the third row is a counterexample to the argument - it shows that the argument is invalid because it is possible for the premise to be T and the conclusion F. If we had some way to know to calculate only this third row, and to ignore the other three, we would have the answer (invalid) with only 1/4 the effort.

So let us see whether we can find a method to locate rows where counterexamples will be. Suppose that the argument  $Q \rightarrow \neg P \vdash P \rightarrow Q$  does have a counterexample. Then there must be a row where  $Q \rightarrow \neg P$  is T and  $P \rightarrow Q$  is F. But there is only one way for a conditional like  $P \rightarrow Q$  to be F, namely when the antecedent (P) is T and the conclusion (Q) is F. This means that if there is a counterexample at all, it must be on the row with  $P = T$  and  $Q = F$ , i.e. the third row. No other rows could possibly count as a counterexample, because on these rows the conclusion will be T. At this point we can easily check that the premise  $Q \rightarrow \neg P$  is indeed T on this row, verifying that row 3 is indeed a counterexample.

Notice how this reasoning focuses our attention on the crucial third row, and helps us see why it is pointless to calculate out the other three rows. By reasoning backwards from the fact that a counterexample requires an F conclusion to the values the letters must have, it may be possible to narrow down a search for a counterexample to one or two rows. (The same idea may also be used on the premises: knowing that a premise has to be T, also yields information on what values the letters have to have in a counterexample.)

You may get the impression that this idea only works for invalid arguments where a counterexample exists. But it works for valid arguments as well. Let us use  $P \rightarrow \neg Q \vdash Q \rightarrow \neg P$  to illustrate. If this argument were to have a counterexample, then there would have to be a row where  $P \rightarrow \neg Q = T$  and  $Q \rightarrow \neg P = F$ . From the latter fact, it follows that the row would have to have  $Q = T$  and  $\neg P = F$ . There is only one row like this, namely where  $Q = T$  and  $P = T$ . But if this row is a counterexample it must make  $P \rightarrow \neg Q = T$ . However, it does not, since  $P = T$  and  $Q = T$  makes  $\neg Q = F$ , with the result that  $P \rightarrow \neg Q = F$ . The only possible counterexample to our argument is the row where  $P = T$  and  $Q = T$ . But we have just seen that it does not make the premise T, so it does not qualify as a counterexample. It follows that the argument has no counterexample, and so it must be valid. Notice that even in the case of this valid argument, we only needed to consider

calculating out values for one of the four rows.

So far we have only explained ideas behind the tree method. It is time to learn the notation and details. The basic idea is to develop a coordinated search for a counterexample to our argument by using the information that the premises must be T and the conclusion F. One notation for doing this involves labeling formulas and their parts with Ts and Fs to indicate their values. However, this can be both conceptually and visually confusing. Instead, the tree method uses the convention that every sentence we write down is considered to be T. If we want to indicate that sentence P is F, we simply write down  $\neg P$  instead, (for if  $\neg P$  is T, P must be F). Let us illustrate this idea by actually building the tree for the argument  $P \rightarrow \neg Q \vdash Q \rightarrow \neg P$ . We begin with the information that if the argument is to have a counterexample in some row, then  $P \rightarrow \neg Q$  must be T and  $Q \rightarrow \neg P$  must be F:

$$\begin{array}{l} P \rightarrow \neg Q \\ \neg(Q \rightarrow \neg P) \end{array}$$

Note that to indicate  $Q \rightarrow \neg P = F$ , we have written  $\neg(Q \rightarrow \neg P)$  instead. Now if  $\neg(Q \rightarrow \neg P)$  is to be T, i.e.  $Q \rightarrow \neg P = F$ , then there is only one possibility:  $Q = T$  and  $\neg P = F$ . Entering this information using our convention, we have the following.

$$\begin{array}{l} P \rightarrow \neg Q \\ \textcircled{1} \neg(Q \rightarrow \neg P) \\ Q \\ \neg\neg P \end{array}$$

The circled 1 is there to indicate that we have worked on the second line - the numbers will indicate the order in which we worked. We have carried out the tree rule for negative arrows. In general, the rule has the form:

$$\begin{array}{l} \neg(a \rightarrow b) \\ a \\ \neg b \end{array}$$

This indicates that whenever we have  $\neg(a \rightarrow b)$  in a tree, we should then write down  $a$  and  $\neg b$  below it. This principle is strongly related to the AR (Arrow) rule. You probably remember the following useful move:  $\neg(a \rightarrow b) \vdash a \& \neg b$ . Notice that the tree rule for a negative arrow amounts to almost the same thing, for from  $\neg(a \rightarrow b)$ , we obtain  $a \& \neg b$  by AR, from which  $a$  and  $\neg b$  follow by  $\&$ Out.

Notice that  $\neg\neg P$  now appears on the tree. Using the double negation rule:

$$\begin{array}{l} \neg\neg a \\ a \end{array}$$


we may simplify this to P:

$$\begin{array}{l} P \rightarrow \neg Q \\ \textcircled{1} \neg(Q \rightarrow \neg P) \\ Q \\ \textcircled{2} \neg\neg P \\ P \end{array}$$


The next step in building our tree corresponds to checking to see whether  $P \rightarrow \neg Q$  can be T. To do so, we must introduce the tree rule for positive conditionals. In general the rule has the following form:

$$\begin{array}{c} a \rightarrow b \\ \swarrow \quad \searrow \\ \neg a \quad b \end{array}$$


The branch in the rule indicates that for  $\alpha \rightarrow \beta$  to be true, there are two (and only two) possibilities: either  $\alpha$  must be F or  $\beta$  must be T. Review the truth table for  $\rightarrow$  to convince yourself that this is correct. Another way to understand (and remember) this rule is to think of it as a variation on the AR rule for converting from  $\vee$  to  $\rightarrow$ :  $\alpha \rightarrow \beta \vdash \neg\alpha \vee \beta$ . When we apply this principle to our tree, we have the following:



The circled numbers show that we have worked on every complex line of our tree. It is now time to evaluate the argument. The tree shows that if there is to be a counterexample to our argument, there are two possibilities. The first, is recorded by the left branch, which contains the simple sentences  $Q$ ,  $P$  and  $\neg P$ . This indicates (according to our convention) that  $Q$  must be T ( $Q$ ),  $P$  must be T ( $P$ ) and  $P$  must be F ( $\neg P$ ). But it is impossible for  $P$  to be both T and F on the same row, so this possibility can be ruled out. To indicate that this situation is impossible, we indicate that this branch is "dead" by adding the contradiction sign:  $\perp$ .



Notice that the right hand branch also contains a contradiction, for the counterexample it describes requires that  $Q$  be both F and T. So we mark that branch with  $\perp$  as well:



Let us review what this tree tells us about validity. It provides a complete record of all the possible ways in which a counterexample can be constructed for  $P \rightarrow \neg Q \vdash Q \rightarrow \neg P$ . There were two options (described by the left and right branches), but both of these turned out to be impossible. The verdict: there cannot be a counterexample to this argument. To put it another way, the argument is valid.

Now we will work out the tree for  $Q \rightarrow \neg P \vdash P \rightarrow Q$  to see what happens with an invalid argument. We begin by writing down the premise and the negative of the conclusion:

$$Q \rightarrow \neg P$$

$$\neg(P \rightarrow Q)$$

Applying the rule for negative conditionals we obtain:

$$Q \rightarrow \neg P$$

$$\textcircled{1} \neg(P \rightarrow Q)$$

$$P$$

$$\neg Q$$

Now we apply the rule for positive conditionals to  $Q \rightarrow \neg P$ :

$$\textcircled{2} Q \rightarrow \neg P$$

$$\textcircled{1} \neg(P \rightarrow Q)$$

$$P$$

$$\neg Q$$

$$\begin{array}{l} \swarrow \quad \searrow \\ \neg Q \quad \neg P \end{array}$$

Note that the right-hand branch contains a contradiction, so we mark it closed.

$$\textcircled{2} Q \rightarrow \neg P$$

$$\textcircled{1} \neg(P \rightarrow Q)$$

$$P$$

$$\neg Q$$

$$\begin{array}{l} \swarrow \quad \searrow \\ \neg Q \quad \neg P \\ \quad \quad \perp \end{array}$$

However, the left hand branch is still open. Since we have worked on every complex sentence, we have a full account of the possibilities for counterexamples for this argument. The left hand branch reports that there is a counterexample, namely one with  $P$ ,  $\neg Q$ , and  $\neg Q$  all true. That amounts to saying that there is a counterexample with  $P = T$  and  $Q = F$ . We list this counterexample by putting the values in a box:

$$\textcircled{2} Q \rightarrow \neg P$$

$$\textcircled{1} \neg(P \rightarrow Q)$$

$$P$$

$$\neg Q$$

$$\text{INVALID}$$

$$\begin{array}{l} \swarrow \quad \searrow \\ \neg Q \quad \neg P \\ \quad \quad \perp \end{array}$$

$P=T$   
 $Q=F$

Here is a review of the tree construction process.

1. Write down the premises and the negative of the conclusion.
2. Apply tree rules to every complex sentence.
3. Mark every branch that contains a contradiction with  $\perp$ .
4. If all branches are closed (contain  $\perp$ ), the argument is valid; if any branch is open (no  $\perp$  on it), then this branch describes a counterexample and the argument is invalid.

Let's review the process with a more complex argument:  $(P \& Q) \rightarrow R$ ,  $P \vdash Q \rightarrow R$ . First enter the premises and the negative of the conclusion:

$$\begin{array}{l} (P \& Q) \rightarrow R \\ P \\ \neg(Q \rightarrow R) \end{array}$$

Now apply the negative conditional rule to the last line:

$$\begin{array}{l} (P \& Q) \rightarrow R \\ P \\ \textcircled{1} \neg(Q \rightarrow R) \\ Q \\ \neg R \end{array}$$

Now apply the positive conditional rule to the first premise:

$$\begin{array}{l} \textcircled{2} (P \& Q) \rightarrow R \\ P \\ \textcircled{1} \neg(Q \rightarrow R) \\ Q \\ \neg R \\ \swarrow \quad \searrow \\ \neg(P \& Q) \quad R \\ \quad \quad \perp \end{array}$$

Notice that the right hand branch is closed because it contains  $\neg R$  and  $R$ , so we have marked it with  $\perp$ . Now we must apply a rule to  $\neg(P \& Q)$ . The rule for negative conjunctions is similar to De Morgan's Law:  $\neg(a \& b) \vdash \neg a \vee \neg b$ . If  $a \& b$  is F then either  $a$  is false or  $b$  is false:


$$\begin{array}{l} \neg(a \& b) \\ \swarrow \quad \searrow \\ \neg a \quad \neg b \end{array}$$

Applying this rule to  $\neg(P \& Q)$ , we obtain:


$$\begin{array}{l} \textcircled{2} (P \& Q) \rightarrow R \\ P \\ \textcircled{1} \neg(Q \rightarrow R) \\ Q \\ \neg R \\ \swarrow \quad \searrow \\ \textcircled{3} \neg(P \& Q) \quad R \\ \swarrow \quad \searrow \quad \perp \\ \neg P \quad \neg Q \quad \text{VALID} \\ \perp \quad \perp \end{array}$$

Since all branches close, the argument is valid.


There are still a few details about the tree method that need explaining. To illustrate, we will work out the tree for  $P \vee Q \vdash P \& Q$ . We begin entering the premise and negative conclusion as usual, and we have already applied the negative  $\&$  rule to the last step:




Now we must work on  $P \vee Q$ . It is easy enough to see what rule we want here. If  $A \vee B$  is T then either  $A$  is T or  $B$  is T:



The problem is where to put the result of applying this rule. We have two open branches in our tree, one ending with  $\neg P$  and the other ending with  $\neg Q$ , and neither one is closed. So where do we place the new branch generated by  $P \vee Q$ :




on the left (below  $\neg P$ ) or on the right (below  $\neg Q$ )? It would be wrong to do either one, and it would also be wrong to divide the results of applying the rule between the two sides like this:



To see what we should do, it helps to remember what the tree represents. The two branches ending with  $\neg P$  and  $\neg Q$  represent two alternatives: for there to be a counterexample, either  $P = F$  or  $Q = F$ . However, the fact that  $P \vee Q$  is T means that on top of these two alternatives there are two more, either  $P = T$  or  $Q = T$ . So the correct thing to do is represent a total of four alternatives. Beneath the alternative  $\neg P$ , we need to record the alternative  $P$  or  $Q$ , and beneath  $\neg Q$  we must do so as well. This illustrates a basic principle about building a tree when there is more than one open branch.

The results of applying a rule to a line must be entered on every open branch below that line.


Following this principle, our tree now looks like this:




Note that contradictions appeared on two of the branches, so we marked them closed: Since we have worked on every step, the tree is complete and we can evaluate the argument. Note that there are two open branches indicating two different counterexamples to our argument. So the argument is invalid. It is a good idea to calculate the truth table for this argument to verify that it has these two counterexamples. One nice thing about trees is that they always give a full account of all the counterexamples to an argument.




Our next project will be to show that  $(P \vee Q) \rightarrow R \vdash (P \rightarrow R) \& (Q \rightarrow R)$  is valid using trees. We begin by negating the conclusion, and applying the negative  $\&$  rule:




Next we will work on the  $\neg(P \rightarrow R)$  on the left hand branch. According to the negative  $\rightarrow$  rule we enter  $P$  and  $\neg R$  below  $\neg(P \rightarrow R)$ . It is not necessary to place  $P$  and  $\neg R$  below the right hand branch. Remember the rule is to put the results of applying a rule on every open branch below the formula you are working on.



After working on  $\neg(Q \rightarrow R)$  on the right branch, we obtain.




Now we will work on the first line:  $(P \vee Q) \rightarrow R$ . In this case, we must place the result of applying the rule on all open branches below this line. So the result goes on both branches:



All that remains is to work on the  $\neg(P \vee Q)$  on each branch. The rule for a negative  $\vee$  is as follows:

$$\begin{array}{c}
 \neg(a \vee b) \\
 \neg a \\
 \neg b
 \end{array}$$

This rule is related to DeMorgan's Law. The idea is that when  $a \vee b$  is F, then both  $a$  and  $b$  are F. Applying this rule on each branch we complete the tree:




There is a final detail about tree construction worth mentioning. You may have been curious about the order in which I have carried out the steps in the trees. You might want to experiment with the problems we have already done to see what happens if you carry out the steps in a different order. In each case I have ordered the steps to keep the tree as simple as possible. There is a simple recipe to help economize in tree construction. The branching rules tend to create open branches, and subsequent steps must be copied onto all these open branches. For this reason, it is a good idea to **do the non-branching steps first**. Then when branching steps are finally carried out, there is a better chance that the branches created will close.


As a last note, we must discuss the  $\leftrightarrow$  rules. The rules may appear complex. But there is a simple way to remember them.



In case  $a \leftrightarrow b$  is T, the values of  $a$  and  $b$  match. So either  $a$  and  $b$  are both T and  $b$  are both F. So the positive  $\leftrightarrow$  rule indicates these two possibilities on two branches. In case  $a \leftrightarrow b$  is F, then the values of  $a$  and  $b$  disagree. So either  $a$  is T and  $b$  is F or  $a$  is F and  $b$  is T. This is why the negative  $\leftrightarrow$  rule shows two alternatives, one containing  $a$  and  $\neg b$  and the other containing  $\neg a$  and  $b$ . Here is a tree for the argument  $M \leftrightarrow G \vdash G \leftrightarrow M$  to help you practice these rules.











Since there are two open branches, the result of applying the negative  $\leftrightarrow$  rule to  $\neg(M \leftrightarrow G)$  goes on both sides:




Note that all branches are closed. It is not necessary for there to be a contradiction for  $M$  and a contradiction for  $G$  to close a branch. One contradiction on a branch is enough to close it.

## Tree Rules for Propositional Logic




Example: To show the argument  $(p \& q) \rightarrow r, p \vdash q \rightarrow r$  is valid, negate the conclusion, and then show that there is no assignment that makes the premises and the negated conclusion all T.



- EXERCISES.
- 1)  $p \rightarrow (q \rightarrow r) \vdash q \rightarrow (p \rightarrow r)$
  - 2)  $p \rightarrow (p \rightarrow q) \vdash p \rightarrow q$
  - 3)  $p \vdash q \vee (r \rightarrow p)$
  - 4)  $p \vdash (p \rightarrow r) \leftrightarrow r$
  - 5)  $p \vee q, p \rightarrow r, q \rightarrow s \vdash r \vee s$

## Finding Counterexamples with Trees

To find a counterexample to this invalid argument:  $p, q \rightarrow \sim p \vdash q$   
 first build its tree:




### Definition

A counterexample is an assignment of truth values to the letters of an argument that makes all premises T and the conclusion F.

The left branch is open (marked with  $\nearrow$ ). Now construct the open branch assignment, by assigning T to all letters that appear unnegated in the open branch, and F to those that appear negated. In this case we have  $p=T$  and  $q=F$ . Calculate the values of the sentences in this argument, and you will find that the premises are T and the conclusion is F in this case.

Here is tree for the argument  $p \rightarrow q, \sim r \rightarrow q \vdash p \rightarrow r$ .




Here is your counterexample for the open branch.

The open branch assignment is  $p=T, q=T, r=F$ . Calculate the values of the sentences in this argument to show that the premises are T and the conclusion is F.


- EXERCISES.
- 1)  $p \rightarrow q \vdash q \rightarrow p$
  - 2)  $p \rightarrow q, p \rightarrow r \vdash \sim r \rightarrow q$
  - 3)  $p \rightarrow q \vdash (p \vee r) \rightarrow q$
  - 4)  $(p \& r) \rightarrow q \vdash p \rightarrow q$

Notes p 25


1. ③  $p \rightarrow (q \rightarrow r)$   
①  $\neg (q \rightarrow (p \rightarrow r))$



2. ②  $p \rightarrow (p \rightarrow q)$




3. P  
①  $\neg (q \vee (r \rightarrow p))$





Notes p. 25




1. ②  $p \rightarrow q$



2. ②  $p \rightarrow q$   
③  $p \rightarrow r$




5. ③  $p \vee q$   
②  $p \rightarrow r$   
④  $q \rightarrow s$   
①  $\neg (r \vee s)$




Ch. 9

3. ②  $F \vee R$   
③  $F \rightarrow S$   
①  $R \rightarrow B$




12. ③  $R \rightarrow F$   
④  $\neg R \rightarrow E$   
①  $(F \vee E) \rightarrow P$




Ch 10


2. ②  $T \vee P$   
①  $\neg T$



4. ①  $C \rightarrow \neg F$   
②  $\neg F$   
③  $\neg C$ , INVALID




3. ①  $S \rightarrow C$   
②  $\neg S$




# Exercises for Trees

## CH 10 (CONTINUED)


- 1c  $D \rightarrow E$   
 ③  $E \vee F$   
 ①  $\neg [(D \vee F) \rightarrow (E \wedge F)]$




5. ④  $P \leftrightarrow W$   
 ③  $\neg (P \leftrightarrow W)$   
 ①  $\neg \neg (P \vee W)$   
 ②  $P \vee W$




6. ①  $E \rightarrow P$   
 ②  $(E \wedge P) \rightarrow T$




10.  $R \vee \neg R$   
 ③  $R \rightarrow (F \wedge S)$   
 ⑤  $\neg R \rightarrow (E \wedge \neg S)$   
 ①  $\neg (F \wedge \neg S)$   
 ②  $F \wedge \neg S$




17. ②  $B \rightarrow W$   
 ③  $\neg B \rightarrow A$   
 ④  $\neg (W \vee A)$




- 19a. ①  $B \rightarrow D$   
 ②  $\neg B \rightarrow D$




8. ⑥  $A \rightarrow C$   
 ⑤  $C \rightarrow A$   
 ②  $A \leftrightarrow C$   
 ①  $\neg [(A \vee C) \rightarrow E]$   
 ④  $A \vee C$



21. ①  $I \vee R$   
 ②  $P \rightarrow I$



24. ②  $U \leftrightarrow (V \wedge F)$   
 ④  $F \rightarrow C$   
 ①  $\neg (C \rightarrow U)$



# Exercise Answers Ch 12

Ch 12 2.  $(S \& R) \& \sim(S \& R)$  is a contradiction

Tree  $(S \& R) \& \sim(S \& R)$  Proof (1)  $S \& R \& \sim(S \& R)$  (PA)  
 $\perp$   
 $\sim(S \& R)$   
 $\perp$   
 $S \& R$   
 $\perp$

4.  $(M \& A) \rightarrow (M \vee A)$  is a logical truth

Tree (1)  $\sim[(M \& A) \rightarrow (M \vee A)]$  Proof (1)  $M \& A$  (PA)  
 (2)  $M \& A$  1 (2)  $M$  1 (2)  $A$   
 (3)  $\sim(M \vee A)$  1 (3)  $M \vee A$  2 (VI)  
 $\perp$   
 $M$   
 $A$   
 $\sim M$   
 $\sim A$   
 $\perp$

7.  $((C \rightarrow D) \rightarrow C) \rightarrow C$  is a logical truth

Tree (1)  $\sim[(C \rightarrow D) \rightarrow C]$  Proof (1)  $(C \rightarrow D) \rightarrow C$  (PA)  
 (2)  $(C \rightarrow D) \rightarrow C$  2 (2)  $\sim C$  (DA)  
 $\perp$   
 $\sim(C \rightarrow D)$  1, 2 MT  
 $C \& \sim D$  3 AR  
 $C$  4 (2)  
 $C \& \sim C$  5 (2) RI  
 $\perp$   
 $C$  2-6  $\rightarrow$  I  
 (8)  $((C \rightarrow D) \rightarrow C) \rightarrow C$  1-7  $\rightarrow$  I

10.  $\sim H \leftrightarrow \sim \sim H$  is a contradiction

Tree (1)  $\sim H \leftrightarrow \sim \sim H$  Proof (1)  $\sim H \leftrightarrow \sim \sim H$  (PA)  
 (2)  $\sim H$  2 (2)  $\sim \sim H$  1 (3)  $\sim H \rightarrow \sim \sim H$  1 (3)  $\rightarrow$  O  
 (3)  $\sim H$  3, 2  $\rightarrow$  O  
 (4)  $\sim H \& \sim \sim H$  4, 2 & I  
 $\perp$   
 $\sim H$  2-5  $\rightarrow$  O  
 (6)  $\sim H$  7 (7)  $\sim \sim H$  1 (8)  $\sim H \rightarrow \sim \sim H$  1 (8)  $\rightarrow$  O  
 (9)  $\sim \sim H$  8, 7  $\rightarrow$  O  
 (10)  $\sim H \& \sim \sim H$  7, 9 & I  
 $\perp$   
 $H$  7-10  $\rightarrow$  O  
 (11)  $H \& \sim H$  11, 6 RI  
 (12)  $\sim H$  1 (12)  $\sim H$   
 (13)  $\sim[\sim H \leftrightarrow \sim \sim H]$  1, 12  $\rightarrow$  I

# Exercise Answers Ch 13

Ch 13 3.  $C \rightarrow S$  and  $C \rightarrow (C \rightarrow S)$  are equivalent

tree

$$C \rightarrow S$$

$$-[C \rightarrow (C \rightarrow S)]$$

$$\begin{array}{l} C \\ -(C \rightarrow S) \\ \perp \end{array}$$

$$C \rightarrow (C \rightarrow S)$$

$$-(C \rightarrow S)$$

$$\begin{array}{l} C \\ -S \\ -C \quad C \rightarrow S \\ \perp \quad \perp \end{array}$$

PROOF

(1)  $C \rightarrow (C \rightarrow S)$

(2)  $C$  (PA)

(3)  $C \rightarrow S$  1, 2  $\rightarrow$  I

(4)  $S$  3, 2  $\rightarrow$  I

(5)  $C \rightarrow S$  2-4  $\rightarrow$  I

(1)  $C \rightarrow S$

(2)  $C$  (PA)

(3)  $C$  (PA)

(4)  $S$  3, 1  $\rightarrow$  I

(5)  $C \rightarrow S$  3-4  $\rightarrow$  I

(6)  $C \rightarrow (C \rightarrow S)$  2-5  $\rightarrow$  I

6.  $(I \vee D) \& A$  and  $I \vee (D \& A)$  are not equivalent

Tree

(1)  $(I \vee D) \& A$

(2)  $-[I \vee (D \& A)]$

(3)  $I \vee D$

$A$

$-I$

$-(D \& A)$

$I$   $D$

$\perp$   $\perp$

(1)  $I \vee (D \& A)$

(2)  $-[ (I \vee D) \& A ]$

$I$   $D \& A$

$D$   $A$

$- (I \vee D)$   $-A$

$-I$   $-D$

$\perp$   $\perp$

$A=F$   
 $I=T$

NO PROOF

9.  $(C \vee D) \rightarrow (-A \& -B)$  and  $(A \vee B) \rightarrow (-C \& -D)$  are equivalent.

TREE

(1)  $(C \vee D) \rightarrow (-A \& -B)$

(2)  $-(A \vee B) \rightarrow (-C \& -D)$

(3)  $A \vee B$

(4)  $-(C \& -D)$

(5)  $-(C \vee D)$

(6)  $-A \& -B$

(7)  $A$   $B$

(8)  $-A$   $-B$

(9)  $C$   $D$

(10)  $-C$   $-D$

(11)  $A$   $B$

(12)  $-A$   $-B$

(13)  $C$   $D$

(14)  $-C$   $-D$

PROOF

(1)  $(A \vee B) \rightarrow (-C \& -D)$

(2)  $C \vee D$

(3)  $-(A \vee B)$

(4)  $-C \& -D$

(5)  $-A$

(6)  $-B$

(7)  $C$   $D$

(8)  $-A$   $-B$

(9)  $C \vee D$   $\&$   $-(C \vee D)$  2, 6  $\&$  I

(10)  $-A \& -B$  3-7  $\rightarrow$  I

(11)  $(C \vee D) \rightarrow (-A \& -B)$  2-8  $\rightarrow$  I

(1)	$C \vee D \rightarrow -A \& -B$	(PA)
(2)	$A \vee B$	
(3)	$-(C \& -D)$	(PA)
(4)	$C \vee D$	3 DM
(5)	$-A \& -B$	1, 4 $\rightarrow$ I
(6)	$-(A \vee B)$	5 DM
(7)	$A \vee B$	2, 6 $\&$ I
(8)	$-C \& -D$	3-7 $\rightarrow$ I
(9)	$A \vee B \rightarrow (-C \& -D)$	2-8 $\rightarrow$ I

(1)	$(A \vee B) \rightarrow (-C \& -D)$	(PA)
(2)	$C \vee D$	
(3)	$-(A \& -B)$	3 DM
(4)	$A \vee B$	1, 4 $\rightarrow$ I
(5)	$-C \& -D$	5 DM
(6)	$-(C \vee D)$	
(7)	$C \vee D$	2, 6 $\&$ I
(8)	$-A \& -B$	3-7 $\rightarrow$ I
(9)	$(C \vee D) \rightarrow (-A \& -B)$	2-8 $\rightarrow$ I



TO S' or	Logical Truth	Contradiction	Consequent	A entails B	A Equiv B to B	more consistent
T. Table	Table for A has all T $\frac{A}{T \quad T}$	Table for A has all F $\frac{A}{F \quad F}$	Table for A has an F and a T $\frac{A}{T \quad F}$	Table for A, B has no T, F row (Argument at B is valid.) $\frac{A \quad B}{T \quad T}$	Tables for A and B are the same $\frac{A \quad B}{T \quad T}$	Table for A, B has a T, T row $\frac{A \quad B}{T \quad T}$
Tree	Tree for $\sim A$ closes $\sim A$ 	Tree for A closes $A$ 	Tree for A open and Tree for ~A open $A$ 	Tree from $A, \sim B$ closes $A$ $\sim B$ 	Trees for both $A, \sim B$ and $B, \sim A$ close $A$ $\sim B$ 	Tree for $A, B$ is open $A$ $B$ 
Proof	Prove A Here's how it looks: $\vdots$ $A$ Goal	Prove $\sim A$ Here's how it looks: $\vdots$ $\sim A$ Goal	No TEST	Assume A, prove B Here's how it looks: $A$ $\vdots$ $B$ Goal	Assume A, prove B Then assume B, and prove A. Here's how it looks: $A$ $\vdots$ $B$ Goal	No TEST

## Predicate Logic Translation Chart

### Singular:

(Use a lower case letter s for the subject and a upper case letter P for the predicate and write: Ps.)

**Examples:** John, David's cat, he\*, she\*, it\*, that\*, the king\*

\* on occasion, these may be general

### General:

#### A. $\forall x(Ax \rightarrow Bx)$

All As are Bs  
Every A is a B  
As are Bs  
Any A is a B  
Each A is a B  
An A is a B\*  
Only Bs are As  
None but Bs are As

\* Usually 'a' means some, but it may mean all. It is NEVER singular.

#### E. $\neg \exists x(Ax \& Bx)$ or $\forall x(Ax \rightarrow \neg Bx)$

No A is a B  
None of the As are Bs  
Not any A is a B

#### I. $\exists x(Ax \& Bx)$

Some As are Bs  
Some A is a B  
There are As that are Bs  
A least one A is a B  
As are sometimes Bs  
An A is a B\*  
Many A are B  
There exist As that are Bs

#### O. $\exists x(Ax \& \neg Bx)$ or $\neg \forall x(Ax \rightarrow Bx)$

Some As are not Bs  
Some As are non-Bs  
Not all As are Bs

# Exercises in Predicatal Logic Ch 2.

1. a Pt      b  $\exists x (Sx \& Nx)$       c  $\exists x (Jx \& -Tx)$       d  $\exists x (Dx \& Sx)$   
 e  $-Th$       f  $\exists x (Hx \& Dx)$       g  $\exists x (Nx \& -Tx)$       h  $\exists x (Gx \& Fx)$   
 i  $\exists x (Px \& -Tx)$       j  $\exists x (Mx \& Fx)$

2. a Janet Reno is a politician.  
 b Janet Reno does not live in Ohio.  
 c Some politicians live in Ohio.  
 d Some politicians live in Ohio.  
 e Some politicians do not live in Ohio.  
 f Some Ohioans are not politicians.

3. a  $\forall x (Lx \rightarrow Bx)$       b  $-\exists x (Wx \& Ex)$  or  $\forall x (Wx \rightarrow -Ex)$       c  $\forall x (Ax \rightarrow Gx)$   
 d Ft      e  $\forall x (Nx \rightarrow Mx)$       f  $\forall x (Wx \rightarrow \sim Ax)$       g  $\forall x (Tx \rightarrow Bx)$   
 h  $-Fs$       i  $\forall x (Sx \rightarrow -Fx)$       j  $\exists x (Px \& Yx)$       k  $\forall x (Ex \rightarrow Cx)$   
 l  $-\exists x (Wx \& Rx)$  or  $\forall x (Wx \rightarrow \sim Rx)$       m  $\forall x (Ix \rightarrow Ex)$       n  $\forall x (Rx \rightarrow Dx)$   
 o  $\exists x (Wx \& \sim Dx)$

4. a  $\forall x (Lx \rightarrow -Cx)$       b  $\forall x (Ix \rightarrow Ax)$       c  $-\forall h$       d  $\forall x (Mx \rightarrow Ux)$   
 e  $\forall x (Rx \rightarrow -Tx)$       f  $\forall x (Kx \rightarrow Lx)$       g  $-\forall x (Px \rightarrow Rx)$  or  $\exists x (Px \& -Rx)$   
 h  $\forall x (Ix \rightarrow Ox)$       i  $-\exists x (Ax \& Rx)$  or  $\forall x (Ax \rightarrow -Rx)$       j  $\forall x (Bx \rightarrow Lx)$   
 k  $\forall x (Dx \rightarrow -Tx)$       l  $\forall x (Sx \rightarrow Lx)$  or  $-\exists x (Sx \& -Lx)$   
 m  $-\exists x (Sx \& Wx)$  or  $\forall x (Sx \rightarrow -Wx)$  or  $\forall x (Wx \rightarrow -Sx)$   
 n  $\forall x (Hx \rightarrow Cx)$

6. a A      b I      c A      d I      e singular      f A

7. a  $\forall x (Px \rightarrow Sx)$       b  $\forall x (Sx \rightarrow Mx)$  or perhaps  $\exists x (Sx \& Mx)$   
 c  $\sim \forall x (Lx \rightarrow Gx)$  or  $\exists x (Lx \& -Gx)$       d  $-\exists x (Mx \& -Px)$  or  $\forall x (Mx \rightarrow Px)$   
 e  $\forall x (Ex \rightarrow Tx)$  'The only A, B' appears to mean 'all A, B'  
 f  $\forall x (Wx \rightarrow Lx)$  or  $-\exists x (Wx \& -Lx)$       g  $-\exists x (Bx \& -Jx)$  or  $\forall x (Bx \rightarrow Jx)$   
 h  $-\forall x (Mx \rightarrow Hx)$  or  $\exists x (Mx \& -Hx)$       i  $\forall x (Wx \rightarrow Cx)$   
 j  $-\exists x (Nx \& -Mx)$  or  $\forall x (Nx \rightarrow Mx)$       k  $\forall x (Wx \rightarrow \sim Sx)$

8. a  $\forall x (Px \rightarrow Ex)$       d  $-\exists x (Px \& Ex)$   
 b  $\forall x (Px \rightarrow Ex)$       e  $\forall x (Px \& Ex) \rightarrow Er$   
 c  $-\forall x (Px \rightarrow Ex)$  or  $\exists x (Px \& -Ex)$       f  $\exists x (Px \& Ex) \rightarrow Er$  or  $\forall x [(Px \& Ex) \rightarrow Er]$

# Exercises in Predicate Logic Ch. 3.1

- a
- |     |                                  |                      |
|-----|----------------------------------|----------------------|
| (1) | $\forall x (Ax \rightarrow -Bx)$ | A                    |
| (2) | BC                               | A                    |
| (3) | $AC \rightarrow -BC$             | 1 $\forall$ D        |
| (4) | $BC \rightarrow -AC$             | 3 CN                 |
| (5) | -AC                              | 4, 2 $\rightarrow$ D |

- b 1
- |      |                                  |                      |
|------|----------------------------------|----------------------|
| (1)  | De                               | A                    |
| (2)  | $\forall x (Dx \rightarrow Fx)$  | A                    |
| (3)  | -Ge                              |                      |
| (4)  | $\forall x (Fx \rightarrow Gx)$  | PA                   |
| (5)  | $De \rightarrow Fe$              | 2 $\forall$ D        |
| (6)  | Fe                               | 5, 1 $\rightarrow$ D |
| (7)  | $Fe \rightarrow Ge$              | 4 $\forall$ D        |
| (8)  | Ge                               | 7, 6 $\rightarrow$ D |
| (9)  | $Ge \wedge -Ge$                  | 8, 3 & I             |
| (10) | $-\forall x (Fx \rightarrow Gx)$ | 4-9 -I               |

2. a
- |     |                                 |                      |
|-----|---------------------------------|----------------------|
| (1) | $\forall x (Sx \rightarrow Ax)$ | A                    |
| (2) | Se                              | A                    |
| (3) | $Se \rightarrow Ae$             | 1 $\forall$ D        |
| (4) | Ae                              | 3, 2 $\rightarrow$ D |

- b
- |     |                                 |                      |
|-----|---------------------------------|----------------------|
| (1) | $\forall x (Ax \rightarrow Fx)$ | A                    |
| (2) | -Fe                             | A                    |
| (3) | $Ae \rightarrow Fe$             | 1 $\forall$ D        |
| (4) | $-Fe \rightarrow -Ae$           | 3 CN                 |
| (5) | -Ae                             | 4, 2 $\rightarrow$ D |

- 3.
- |     |                                  |                      |
|-----|----------------------------------|----------------------|
| (1) | $\forall x (Rx \rightarrow -Ox)$ | A                    |
| (2) | Ra                               | A                    |
| (3) | $Ra \rightarrow -Oa$             | 1 $\forall$ D        |
| (4) | -Oa                              | 3, 2 $\rightarrow$ D |

- c 1
- |     |                                  |                      |
|-----|----------------------------------|----------------------|
| (1) | Sm                               | A                    |
| (2) | -Fm                              | A                    |
| (3) | $\forall x (Sx \rightarrow Fx)$  | PA                   |
| (4) | $Sm \rightarrow Fm$              | 3 $\forall$ D        |
| (5) | Fm                               | 4, 1 $\rightarrow$ D |
| (6) | $Fm \wedge -Fm$                  | 5, 2 & I             |
| (7) | $-\forall x (Sx \rightarrow Fx)$ | 3-6 -I               |

- 4.
- |     |                                  |                      |
|-----|----------------------------------|----------------------|
| (1) | $\forall x (Lx \rightarrow Ax)$  | A                    |
| (2) | $\forall x (Ax \rightarrow -Mx)$ | A                    |
| (3) | Mt                               | A                    |
| (4) | $At \rightarrow -Mt$             | 2 $\forall$ D        |
| (5) | $Mt \rightarrow -At$             | 4 CN                 |
| (6) | -At                              | 5, 3 $\rightarrow$ D |
| (7) | $Lt \rightarrow At$              | 1 $\forall$ D        |
| (8) | $-At \rightarrow -Lt$            | 7 CN                 |
| (9) | -Lt                              | 8, 6 $\rightarrow$ D |

- 5.
- |     |                                 |                      |
|-----|---------------------------------|----------------------|
| (1) | $\forall x (Px \rightarrow Ax)$ | A                    |
| (2) | $\forall x (Ax \rightarrow Mx)$ | A                    |
| (3) | -Ms                             | A                    |
| (4) | $As \rightarrow Ms$             | 2 $\forall$ D        |
| (5) | $-Ms \rightarrow -As$           | 4 CN                 |
| (6) | -As                             | 5, 3 $\rightarrow$ D |
| (7) | $Ps \rightarrow As$             | 1 $\forall$ D        |
| (8) | $-As \rightarrow -Ps$           | 7 CN                 |
| (9) | -Ps                             | 8, 6 $\rightarrow$ D |

6. 1
- |     |                                   |                      |
|-----|-----------------------------------|----------------------|
| (1) | Ws                                | A                    |
| (2) | Hs                                | A                    |
| (3) | $\forall x (Hx \rightarrow -Wx)$  | PA                   |
| (4) | $Hs \rightarrow -Ws$              | 3 $\forall$ D        |
| (5) | -Ws                               | 4, 2 $\rightarrow$ D |
| (6) | $Ws \wedge -Ws$                   | 1, 5 & I             |
| (7) | $-\forall x (Hx \rightarrow -Wx)$ | -In                  |

# Exercises in Predicate Logic Ch 3, 2

- 8.1 (1)
- 2 (2)
- 3 (3)
- 2 (4)
- 1 (5)
- 3 (6)
- 2 (7)
- 1,2 (8)
- 2 (9)
- 1,3 (10)
- 1,2,3 (11)
- 1,2 (12)

$\forall x (Ax \rightarrow Bx)$	A
$\exists x (Ax \wedge Cx)$	A
$\forall x (Cx \rightarrow -Bx)$	PA
$Aa \wedge Ca$	2 $\exists O$
$Aa \rightarrow Bd$	1 $\forall O$
$Cd \rightarrow -Bd$	3 $\forall O$
$Ad$	4 $\& O$
$Bd$	5,7 $\rightarrow O$
$Cd$	4 $\& O$
$-Bd$	6,9 $\rightarrow O$
$Bd \wedge -Bd$	8,10 $\& I$
$-\forall x (Cx \rightarrow -Bx)$	3-11 -I

- 10.1 (1)
- 2 (2)
- 3 (3)
- 3 (4)
- 1 (5)
- 2 (6)
- 3 (7)
- 1,3 (8)
- 1,2,3 (9)
- 3 (10)
- 1,2,3 (11)
- 1,2 (12)

$\forall x (Wx \rightarrow Fx)$	A
$\forall x (Gx \rightarrow -Cx)$	A
$\exists x (Wx \wedge Cx)$	PA
$Wa \wedge Ca$	3 $\exists O$
$Wa \rightarrow Ga$	1 $\forall O$
$Ga \rightarrow -Ca$	2 $\forall O$
$Wa$	4 $\& O$
$Ga$	5,7 $\rightarrow O$
$-Ca$	6,8 $\rightarrow O$
$Ca$	4 $\& O$
$Ca \wedge -Ca$	10,9 $\& I$
$-\exists x (Wx \wedge Cx)$	3-11 -I

- 11. 1 (1)
- 2 (2)
- 3 (3)
- 1 (4)
- 2 (5)
- 3 (6)
- 1 (7)
- 1,2 (8)
- 1 (9)
- 3 (10)
- 1,2,3 (11)
- 1,2 (12)

$\exists x (Nx \wedge Cx)$	A
$\forall x (Nx \rightarrow -Fx)$	A
$\forall x (Cx \rightarrow Fx)$	PA
$Na \wedge Ca$	1 $\exists O$
$Na \rightarrow -Fa$	2 $\forall O$
$Ca \rightarrow Fa$	3 $\forall O$
$Na$	4 $\& O$
$-Fa$	5,7 $\rightarrow O$
$Ca$	4 $\& O$
$Fa$	6,9 $\rightarrow O$
$Fa \wedge -Fa$	10,8 $\& I$
$-\forall x (Cx \rightarrow Fx)$	3-11 -I















9. 1,9 (1)	$\forall x (Ox \rightarrow Dx)$	A
2 (2)	$\exists x (Ox \wedge -Dx)$	PA
2 (3)	$Oa \wedge -Da$	2 $\exists O$
1 (4)	$Oa \rightarrow Da$	1 $\forall O$
2 (5)	$Da$	3 $\& O$
1,2 (6)	$Da$	4,5 $\rightarrow O$
2 (7)	$-Da$	3 $\& O$
1,2 (8)	$Da \wedge -Da$	4,7 $\& I$
1 (9)	$-\exists x (Ox \wedge -Dx)$	2-8 -I

10. 1 (1)	$\forall x (Bx \rightarrow -Ax)$	A
2 (2)	$\exists x (Bx \wedge Ax)$	PA
2 (3)	$Ba \wedge Aa$	2 $\exists O$
1 (4)	$Ba \rightarrow -Aa$	1 $\forall O$
2 (5)	$Ba$	3 $\& O$
1,2 (6)	$-Aa$	4,5 $\rightarrow O$
2 (7)	$Aa$	3 $\& O$
1,2 (8)	$Aa \wedge -Aa$	7,6 $\& I$
1 (9)	$-\exists x (Bx \wedge Ax)$	2-8 -I

12. 1 (1)	$\forall x (Cx \rightarrow Tx)$	A
2 (2)	$\exists x (Cx \wedge -Dx)$	A
3 (3)	$\forall x (Tx \rightarrow Dx)$	PA
2 (4)	$Ca \wedge -Da$	2 $\exists O$
1 (5)	$Ca \rightarrow Ta$	1 $\forall O$
3 (6)	$Ta \rightarrow Da$	3 $\forall O$
2 (7)	$Ca$	4 $\& O$
1,2 (8)	$Ta$	5,7 $\rightarrow O$
1,2,3 (9)	$Da$	6,8 $\rightarrow O$
2 (10)	$-Da$	4 $\& O$
1,2,3 (11)	$Da \wedge -Da$	9,10 $\& I$
1,2 (12)	$-\forall x (Tx \rightarrow Dx)$	3-11 -I

# Trees for Exercises in Ch 3 of Predicate Logic

n3



# Exercises in Predicate Logic

## Ch 4.2

10. a

(1)	$\neg \exists x (Ox \& \neg Dx)$	A
(2)	$\neg \forall x (Ox \rightarrow Dx)$	PA
(3)	$\forall x \neg (Ox \& \neg Dx)$	1 QE
(4)	$\exists x \neg (Ox \rightarrow Dx)$	2 QE
(5)	$\neg (Oe \rightarrow De)$	4 FO
(6)	$\neg (Oe \& \neg De)$	3 VO
(7)	$Oe \rightarrow De$	6 AR
(8)	$(Oe \rightarrow De) \& \neg (Oe \rightarrow De)$	7, 5 & I
(9)	$\forall x (Ox \rightarrow Dx)$	2-8 -O

b

(1)	$\neg \exists x (Bx \& \neg Ax)$	A
(2)	$\neg \forall x (Bx \rightarrow \neg Ax)$	PA
(3)	$\forall x \neg (Bx \& \neg Ax)$	1 QE
(4)	$\exists x \neg (Bx \rightarrow \neg Ax)$	2 QE
(5)	$\neg (Bf \rightarrow \neg Af)$	4 FO
(6)	$\neg (Bf \& \neg Af)$	3 VO
(7)	$\neg Bf \vee \neg \neg Af$	6 DM*
(8)	$Bf \rightarrow \neg \neg Af$	7 AR
(9)	$(Bf \rightarrow \neg \neg Af) \& \neg (Bf \rightarrow \neg \neg Af)$	8, 5 & I
(10)	$\forall x (Bx \rightarrow \neg \neg Ax)$	2-9 -I

\* I will accept 6 AR here without the DM step.

11. a

(1)	$\forall x (Bx \rightarrow Dx)$	A
(2)	$\forall x (Mx \rightarrow \neg Dx)$	A
(3)	$\neg \forall x (Mx \rightarrow \neg Bx)$	PA
(4)	$\exists x \neg (Mx \rightarrow \neg Bx)$	3 QE
(5)	$\neg (Ma \rightarrow \neg Ba)$	4 FO
(6)	$Ba \rightarrow Da$	1 VO
(7)	$Ma \rightarrow \neg Da$	2 VO
(8)	$Ma \& Ba$	5 AR
(9)	$Ma$	8 & O
(10)	$\neg Da$	7, 9 $\rightarrow$ O
(11)	$Ba$	8 & O
(12)	$Da$	6, 11 $\rightarrow$ O
(13)	$Da \& \neg Da$	12, 10 & I
(14)	$\forall x (Mx \rightarrow \neg Bx)$	3-13 -O

12. a

(1)	$\forall x (Px \rightarrow Sx)$	A
(2)	$\forall x (Sx \rightarrow Rx)$	A
(3)	$\neg \forall x (Px \rightarrow Rx)$	PA
(4)	$\exists x \neg (Px \rightarrow Rx)$	3 QE
(5)	$\neg (Pa \rightarrow Ra)$	4 FO
(6)	$Pa \rightarrow Sa$	1 VO
(7)	$Sa \rightarrow Ra$	2 VO
(8)	$Pa \& \neg Ra$	5 AR
(9)	$Pa$	8 & O
(10)	$Sa$	6, 9 $\rightarrow$ O
(11)	$Ra$	7, 10 $\rightarrow$ O
(12)	$\neg Ra$	8 & O
(13)	$Ra \& \neg Ra$	11, 12 & I
(14)	$\forall x (Px \rightarrow Rx)$	3-13 -O

3. a

(1)	$Wh$	A
(2)	$Fh$	A
(3)	$\neg \exists x (Fx \& Wx)$	PA
(4)	$\forall x \neg (Fx \& Wx)$	3 QE
(5)	$\neg (Fh \& Wh)$	4 VO
(6)	$\neg Fh \vee \neg Wh$	5 DM
(7)	$Fh \rightarrow \neg Wh$	6 AR*
(8)	$\neg Wh$	7, 2 $\rightarrow$ O
(9)	$Wh \& \neg Wh$	8, 1 & I
(10)	$\exists x (Fx \& Wx)$	3-9 -O

12. b

(1)	$\forall x (Lx \rightarrow Ax)$	A
(2)	$\forall x (Ax \rightarrow \neg Rx)$	A
(3)	$\neg \forall x (Lx \rightarrow \neg Rx)$	PA
(4)	$\exists x \neg (Lx \rightarrow \neg Rx)$	3 QE
(5)	$\neg (Lg \rightarrow \neg Rg)$	4 FO
(6)	$Lg \rightarrow \neg Rg$	1 VO
(7)	$Ag \rightarrow \neg Rg$	2 VO
(8)	$Lg \& Rg$	5 AR
(9)	$Lg$	8 & O
(10)	$Ag$	6, 9 $\rightarrow$ O
(11)	$\neg Rg$	7, 10 $\rightarrow$ O
(12)	$Rg$	8 & O
(13)	$Rg \& \neg Rg$	12, 11 & I
(14)	$\forall x (Lx \rightarrow \neg Rx)$	3-13 -O

will also accept 5 AR here without the DM step.

13c	(1) $\exists x (Fx \& Vx)$	A
	(2) $\forall x (Ex \rightarrow -Fx)$	A
2	(3) $-\exists x (Vx \& -Ex)$	PA
3	(4) $\forall x -(Vx \& -Ex)$	3 QE
3	(5) $Fa \& Va$	1 $\exists O$
1	(6) $Ea \rightarrow -Fa$	2 $\forall O$
2	(7) $-(Va \& -Ea)$	4 $\forall O$
3	(8) $Va \rightarrow Ea$	7 AR
3	(9) $Va$	5 $\& O$
1	(10) $Ea$	8, 9 $\rightarrow O$
1, 3	(11) $-Fa$	6, 10 $\rightarrow O$
1, 2, 3	(12) $Fa$	5 $\& O$
1	(13) $Fab - Fa$	12, 11 $\& I$
1, 2, 3	(14) $\exists x (Vx \& -Ex)$	3-13 - O

13d	(1) $\forall x (Bx \rightarrow Mx)$	A
1	(2) $\exists x (Ex \& Bx)$	A
2	(3) $-\exists x (Ex \& Mx)$	PA
3	(4) $\forall x -(Ex \& Mx)$	3 QE
3	(5) $Eb \& Bb$	2 $\exists O$
2	(6) $Bb \rightarrow Mb$	1 $\forall O$
1	(7) $-(Eb \& Mb)$	4 $\forall O$
3	(8) $Bb$	5 $\& O$
2	(9) $Mb$	6, 8 $\rightarrow O$
4, 2	(10) $Eb$	5 $\& O$
2	(11) $Eb \& Mb$	10, 9 $\& I$
1, 2, 3	(12) $(Eb \& Mb) \& -(Eb \& Mb)$	11, 7 $\& I$
1, 2	(13) $\exists x (Ex \& Mx)$	3-12 - O

14	(1) $\forall x (Px \rightarrow Fx)$	A
1	(2) $\forall x (Mx \rightarrow Jx)$	A
2	(3) $\forall x (Fx \rightarrow -Jx)$	A
3	(4) $-\forall x (Mx \rightarrow -Px)$	PA
4	(5) $\exists x -(Mx \rightarrow -Px)$	4 QE
4	(6) $-(Mc \rightarrow -Pc)$	5 $\exists O$
4	(7) $Pc \rightarrow Fc$	1 $\forall O$
1	(8) $Mc \rightarrow Jc$	2 $\forall O$
2	(9) $Fc \rightarrow -Jc$	3 $\forall O$
3	(10) $Mc \& Pc$	6 AR
4	(11) $Mc$	10 $\& O$
4	(12) $Jc$	8, 11 $\rightarrow O$
4	(13) $Pc$	10 $\& O$
4	(14) $Fc$	7, 13 $\rightarrow O$
4	(15) $-Jc$	9, 14 $\rightarrow O$
4	(16) $Jc \& -Jc$	12, 15 $\& I$
1, 2, 3, 4	(17) $\forall x (Mx \rightarrow -Px)$	4-16 - O


15	(1) $\forall x (Tx \rightarrow Px)$	A
1	(2) $\forall x (Dx \rightarrow Ux)$	A
2	(3) $\forall x (Ux \rightarrow Nx)$	A
3	(4) $-\forall x (Tx \rightarrow Nx)$	PA
4	(5) $\exists x -(Tx \rightarrow Nx)$	4 QE
4	(6) $-(Ta \rightarrow Na)$	5 $\exists O$
4	(7) $Ta \rightarrow Da$	1 $\forall O$
1	(8) $Da \rightarrow Ua$	2 $\forall O$
2	(9) $Ua \rightarrow Na$	3 $\forall O$
3	(10) $Ta \& -Na$	6 AR
4	(11) $Ta$	10 $\& O$
4	(12) $Da$	7, 11 $\rightarrow O$
1, 4	(13) $Ua$	8, 12 $\rightarrow O$
1, 2, 4	(14) $Na$	9, 13 $\rightarrow O$
4	(15) $-Na$	6 $\& O$
1, 2, 3, 4	(16) $Na \& -Na$	14, 15 $\& I$
1, 2, 3	(17) $\forall x (Tx \rightarrow Nx)$	4-16 - O

\* See Appendix 4 p 248 for a shorter solution




Trees for trees in Ch 4 of Practical Logic


- 10a
- ①  $\neg \exists x (Ox \& \neg Dx)$
  - ②  $\neg \forall x (Ox \rightarrow Dx)$
  - ③  $\forall x \neg (Ox \& \neg Dx)$
  - ④  $\exists x \neg (Ox \rightarrow Dx)$
  - ⑤  $\neg (Ode \rightarrow De)$
  - ⑥  $\neg (Ode \& \neg De)$




- 10b
- ①  $\neg \exists x (Bx \& Ax)$
  - ②  $\neg \forall x (Bx \rightarrow \neg Ax)$
  - ③  $\forall x \neg (Bx \& Ax)$
  - ④  $\exists x \neg (Bx \rightarrow \neg Ax)$
  - ⑤  $\neg (Bf \rightarrow \neg Af)$
  - ⑥  $\neg (Bf \& \neg Af)$




- 11
- ③  $\forall x (Bx \rightarrow Dx)$
  - ④  $\forall x (Mx \rightarrow \neg Dx)$
  - ①  $\neg \forall x (Mx \rightarrow \neg Bx)$
  - ②  $\exists x \neg (Mx \rightarrow \neg Bx)$
  - ③  $\neg (Ma \rightarrow \neg Ba)$
  - ④  $Ba \rightarrow Da$
  - ⑤  $Ma \rightarrow \neg Da$




- 2a
- ⑤  $\forall x (Px \rightarrow Sx)$
  - ④  $\forall x (Sx \rightarrow Rx)$
  - ①  $\neg \forall x (Px \rightarrow Rx)$
  - ②  $\exists x \neg (Px \rightarrow Rx)$
  - ③  $\neg (Pb \rightarrow Rb)$
  - ④  $Pb \rightarrow Sb$
  - ⑤  $Sb \rightarrow Rb$




- 12b
- ③  $\forall x (Lx \rightarrow Ax)$
  - ④  $\forall x (Ax \rightarrow \neg Rx)$
  - ①  $\neg \forall x (Lx \rightarrow \neg Rx)$
  - ②  $\exists x \neg (Lx \rightarrow \neg Rx)$
  - ③  $\neg (Lg \rightarrow \neg Rg)$
  - ④  $Lg \rightarrow Ag$
  - ⑤  $Ag \rightarrow \neg Rg$




- 13a
- wh
  - Fh
  - ①  $\neg \exists x (Fx \& Wx)$
  - ②  $\forall x \neg (Fx \& Wx)$
  - ③  $\neg (Fh \& Wh)$




- 13b
- ③  $\forall x (Mx \rightarrow Dx)$
  - ④  $\exists x (Mx \& \neg Sx)$
  - ①  $\neg \exists x (Dx \& \neg Sx)$
  - ②  $\forall x \neg (Dx \& \neg Sx)$
  - ③  $Ma \& \neg Sa$
  - ④  $Ma \rightarrow Da$
  - ⑤  $\neg (Da \& \neg Sa)$




- 13c
- ②  $\exists x (Fx \& \forall x)$
  - ③  $\forall x (Ex \rightarrow \neg Fx)$
  - ①  $\neg \exists x (Ex \& \neg Fx)$
  - ④  $\forall x \neg (Ex \& \neg Fx)$
  - ⑤  $Fa \& \forall a$
  - ⑥  $Ea \rightarrow \neg Fa$
  - ⑦  $\neg (\forall a \& \neg Ea)$



- 13d
- ③  $\forall x (Bx \rightarrow Mx)$
  - ④  $\exists x (Ex \& Bx)$
  - ①  $\neg \exists x (Ex \& Mx)$
  - ②  $\forall x \neg (Ex \& Mx)$
  - ③  $Eb \& Bb$
  - ④  $Bb \rightarrow Mb$
  - ⑤  $\neg (Eb \& Mb)$



- 14
- ③  $\forall x (Px \rightarrow Fx)$
  - ④  $\forall x (Mx \rightarrow \neg Fx)$
  - ⑤  $\forall x (Fx \rightarrow \neg Mx)$
  - ①  $\neg \forall x (Mx \rightarrow \neg Px)$
  - ②  $\exists x \neg (Mx \rightarrow \neg Px)$
  - ③  $\neg (Mc \rightarrow \neg Pc)$
  - ④  $Pc \rightarrow Fc$
  - ⑤  $Mc \rightarrow \neg Ic$
  - ⑥  $Fc \rightarrow \neg Ic$



- 15
- ③  $\forall x (Tx \rightarrow Dx)$
  - ④  $\forall x (Dx \rightarrow Ux)$
  - ⑤  $\forall x (Ux \rightarrow Nx)$
  - ①  $\neg \forall x (Tx \rightarrow Nx)$
  - ②  $\exists x \neg (Tx \rightarrow Nx)$
  - ③  $\neg (Ta \rightarrow Na)$
  - ④  $Ta \rightarrow Da$
  - ⑤  $Da \rightarrow Ua$
  - ⑥  $Ua \rightarrow Na$

