
Socio-Demographic Factors Affecting Knowledge Level of Tuberculosis Patients in Rajshahi City, Bangladesh

Nazrul Islam Mondal

Department of Population Science and
Human Resource Development
University of Rajshahi, Bangladesh

Nazrul Hoque

The Hobby Center for Public Policy
University of Houston

Rocky Khan Chowdhury

Department of Population Science
and Human Resource Development
University of Rajshahi, Bangladesh

Jeffrey Howard

Department of Demography
University of Texas at San Antonio

Socio-Demographic Factors Affecting Knowledge Level of Tuberculosis Patients in Rajshahi City, Bangladesh*

Nazrul Islam Mondal, *University of Rajshahi*

Nazrul Hoque, *University of Houston*

Rocky Khan Chowdhury, *University of Rajshahi*

Jeffrey Howard, *University of Texas at San Antonio*

Every year almost two million people die worldwide due to Tuberculosis (TB) and most deaths occur in low- and middle-income countries. In 2010, Bangladesh ranked sixth among 22 high burden countries (HBCs) where, the incidence rate was 225 and mortality rate was 43 per 100,000 population for TB. However, Tuberculosis control program in Bangladesh is still unsatisfactory due to insufficient knowledge and stigma attached to TB. Patients with low knowledge may be at higher risk of experiencing delays in diagnosis and appropriate treatment. The aims of this study were to identify the knowledge levels of TB and investigate the factors associated with knowledge level among the TB patients in Bangladesh. A cross-sectional study was conducted at Rajshahi City, Bangladesh. A total of 384 TB patients were interviewed through a pretested, structured questionnaire using purposive sampling techniques. Logistic regression analysis was used to evaluate the effects of selected socio-demographic factors on TB knowledge level. The results revealed that pulmonary TB patients had greater knowledge than that of extra-pulmonary patients, and that sex, age, educational status and TB type were significantly associated with knowledge level. In general, males and young adults, ages 21-35, had greater awareness about transmission and prevention of TB than females and adults over 35. Individuals with higher education and urban area patients were comparatively better informed about TB infection. Patients with greater knowledge about TB were also less likely to experience delays in seeking treatment.

Keywords: Tuberculosis, National TB program, Rajshahi City, Knowledge index, Logistic regression model.

Introduction

Tuberculosis (TB) is a chronic communicable bacterial disease that remains an important public health problem, especially in developing countries. TB is an airborne, infectious disease caused by bacteria which primarily affect the lungs. Approximately one third of the world's population carries the TB bacteria namely *Mycobacterium TB* (MTB). The World Health Organization (WHO) declared TB as a 'global emergence' in 1993 (Gupta et al. 2002). Every year almost two million people die worldwide due to TB and most deaths occur in low- and middle-income countries (World Health Organization [WHO] 2011). Although TB is a curable disease, it ranks as the second leading cause of death among infectious diseases worldwide, after the human immunodeficiency virus (HIV). TB takes advantage of individual's with weakened immune systems, which is why it is called an opportunistic infectious disease.

* An earlier version of this paper has been published in the Journal of *African Health Sciences*.

Consequently, the risk of TB infection is higher among the people who are HIV positive (Mondal and Shitan 2013a, 2013b). Among 22 high burden countries (HBCs) Bangladesh has been ranked 6th where, the incidence rate for TB was 225 per 100,000 population and TB mortality rate was 43 per 100,000 population in 2010 (WHO 2011). To fight against TB, the Bangladesh National TB Control Program (NTP) has adopted the directly observed treatment short course (DOTs) strategy since 1993 (Zafar Ullah et al. 2006). At present, Bangladesh has more than 165 million people, and is the seventh most populous country in the world. It is also one of the poorest nations, and faces great challenges in providing health care services including TB services for its citizens. People having symptoms of TB should be identified when they seek care at a general health facility, and referred to the specialized TB health care centers for diagnosis, treatment and case management. Given the challenges facing Bangladesh's health services infrastructure, this is often a difficult goal to achieve.

Early case detection depends on patients' perception about their needs of seeking healthcare. Consequently, it is very important to make people understand when and where they should seek healthcare. Health knowledge allows individuals to assess symptoms, identify causes and transmission routes, and provide familiarity with the availability of treatment and cure. Likewise, knowledge and awareness of TB is very important among TB affected people. Increasing knowledge will lead to overcome some of the challenges to control TB. While people may have a general idea of what TB is and how it is treated, gaps in knowledge, such as transmission, treatment, and prevention causes diagnostic and treatment delays among many people living with TB. Delays in treatment occur for several reasons, such as, lack of knowledge, lack of awareness of the significance of symptoms, negative social attitudes or different combinations of these three factors (Koay 2004). Patients with low knowledge about symptoms are less likely to seek healthcare and get diagnosed. Patients with low knowledge are more likely to visit traditional healers and pharmacists rather than DOTs providers, which leads to delays in diagnosis and appropriate treatment.

Although most deaths due to TB occurred among men, the burden of TB is high among women as well. In 2012, an estimated 410,000 women died from TB. In Bangladesh, women tend to have longer diagnostic and treatment delays compared to men (Karim et al. 2007). Lack of TB knowledge has been shown to be associated with diagnostic delay and in some case it is associated with poverty (Ngadaya et al. 2009; Long and Wang 2008; Demissie et al. 2002; Mauch et al. 2011). Bangladesh is considered to be a low human development country based on the value of Human Development Index (HDI=0.500), placed 146 out of 187 countries and territories (UNDP 2011). Consequently, Bangladesh has not had much success in the areas of education and health. Obviously, education is significantly associated with health and makes a great impact on the enhancement on people's knowledge about TB (Wang et al. 2008). To-date, no study has been published with a focus on knowledge level and socio-demographic factors among TB patients in Rajshahi City, Bangladesh. Therefore, the purposes of the study were to identify the knowledge level of TB patients, and to investigate the socio-demographic factors associated with knowledge level.

Background

In this study, the World Health Organization's (WHO) standard definition of TB was followed. TB cases are defined as individuals diagnosed with TB by a health worker or other medical practitioner who has decided to treat the patient with a full course of anti-TB treatment. A pulmonary TB patient is defined as a patient who has TB, which refers to a disease involving the lung parenchyma. An extra-pulmonary TB patient is defined as a patient who has TB


disease, but refers to TB of organs other than the lungs, e.g. pleura, lymph nodes, abdomen, genitourinary tract, skin, joints and bones, meninges, central nervous system, spine, kidneys, pericardium, intestines and peritoneum. A new case is defined by the Bangladesh NTP as being a patient who had never received anti-TB treatment or who had received it for less than 1 month after diagnosis by the government or non-government medical providers. An old case is defined as a patient who has been declared cured but remains a TB patient or not cured after completing continuous phase.

Materials and Methods

Study Area

A cross sectional study was conducted in Rajshahi City, Bangladesh. Bangladesh is administratively divided into seven divisions and Rajshahi is one of them. Rajshahi is the third most populous division with more than 21 million people, accounting for almost 13 percent of the Country's total population. The study area is located in the eastern part of the country, and was selected randomly. The study was carried out in the six different healthcare centers located in the city, namely Rajshahi Medical Collage Hospital (RMCH), Tilottoma (Noudapara Branch), Tilottoma (Bulonpur Branch), Rajshahi Chest Disease Hospital (CDH), Rajshahi Chest Disease Clinic (CDC) and Population Service and Training Centre (PSTC) (Figure 1). These study areas provide free TB treatment under the directly observed therapy (DOT) program, which aims to control TB. The TB control program is a national program of the Directorate General of Health Services (DGHS) under the Mycobacterium Disease Control (MBDC) unit which is run through National TB Control Programme (NTP).

Figure 1 Study Areas



Respondents in this study include patients who began to receive anti-TB treatment under DOT from June 2011 to February 2012. Patients in whom TB is suspected are sent to the laboratory for sputum microscopy and are registered in the TB laboratory register. Patients diagnosed with smear-positive TB are registered for treatment in the TB treatment program. Smear-positive pulmonary TB is diagnosed using direct sputum microscopy examination at the study areas, while smear-negative pulmonary and extra-pulmonary TB is diagnosed by a graduate physician at hospital and respective clinic. After diagnosis, most TB cases are treated and managed by study areas. The study population consisted of both pulmonary and extra-pulmonary TB cases, and further divided into new and old cases, which are identified in the study sites by other diagnostic tools.

Data Collection

Three hundred and eighty four TB patients were interviewed face-to-face through a structured questionnaire using purposive sampling techniques. Information regarding knowledge related to TB (Table 1) and socio-demographic data (Table 3) were collected from 384 respondents. Two trained interviewers and a medical doctor conducted the interviews after obtaining informed consent at the study sites. Patients were informed clearly about the purpose of study. Both types of patients, pulmonary and extra-pulmonary were receiving treatment at the study sites. Pulmonary sputum-positive TB was identified on the basis of at least 1 positive sputum culture of MTB or 2 sputum smears containing acid fast bacilli in the context of a compatible clinical illness. Pulmonary sputum-negative TB was identified on the basis of negative smears and cultures for MTB in the context of clinically and radiologically compatible illness. Diagnosis of extra-pulmonary TB was based on a combination of clinical, radiological, and histopathological findings. For each patient, the following socio-demographic information was collected: sex, age, educational status; residence, delay in seeking treatment, TB type; and knowledge regarding symptoms, transmission, treatment and prevention of TB.

Data Management and Statistical Analysis

Data entry was performed using EPI info software and then was exported to the statistical package (SPSS 16.0) to analyze the data. Univariate analysis was completed to find the distributions of different dimensions of knowledge items and some selected socio-demographic characteristics. Indices of each of the four dimensions of knowledge concerning symptoms (DKS), transmission (DKT), treatment (DKTr), and prevention (DKP) were constructed using the sums of weighted binary input variables where maximum and minimum values were chosen for each underlying dimension. Performance in each dimension is expressed as a unit-free index between 0 and 1 in accordance with the construction method of the Human development Index [13] using the following equation:

$$\text{Dimension Index (DI)} = \frac{\text{Actual value} - \text{Minimum value}}{\text{Maximum value} - \text{Minimum value}}.$$

Based on the above equation, dimension scores for symptoms were computed as follows;
Dimension Index for Symptoms (DIKS) =

$$\frac{\text{Actual value of symptoms} - \text{Minimum value of symptoms}}{\text{Maximum value of symptoms} - \text{Minimum value of symptoms}}.$$

Scores for each of the indices were averaged in order to compute the overall knowledge index (OKI) recorded as:

$$\text{OKI} = \frac{\text{DIKS} + \text{DIKT} + \text{DIKTr} + \text{DIKP}}{4}.$$

The scores obtained for this index were then transformed into a dichotomous variable with categories labeled low/moderate knowledge and sufficient knowledge. Cronbach's α coefficient was used to evaluate the internal reliability of the OKI. The calculated value of α 0.785, suggesting good internal consistency. The Pearson's Chi-squared (χ^2) test was used to determine associations between OKI and socio-demographic predictors. Logistic regression analysis was used to determine the effects of selected socio-demographic predictors on knowledge level.

Explanatory Variables

The dependent variable in our analysis is the knowledge level of the TB patients. The knowledge related variables were composed of four items. These items include knowledge about symptoms, knowledge about transmission, knowledge about treatment, and knowledge about prevention. Knowledge about symptoms includes, coughing with and without blood more than 3 weeks, chest pain; shortness of breath, loss of appetite, weight loss and fever with night sweating. Knowledge about transmission includes, understanding whether TB is communicable through sneezing, through air and through touching items from affected people. Knowledge about treatment includes understanding that treatment is available, is treatment free, can regular intake of medicine cure a patient and does irregular intake of medicine cause death. Knowledge about prevention includes understanding that vaccine is available, stay far away from affected people when they sneeze, do not use items from affected people and always stay clean. The categories of the aforesaid questions are shown in Table 1. The dependent variable, knowledge level, in this study was categorized in two groups: low/moderate knowledge and sufficient knowledge. The dependent variable is considered for logistic regression model is coded in the following way:

$$y = \begin{cases} 1, & \text{is the knowledge level sufficient;} \\ 0, & \text{otherwise.} \end{cases}$$

A number of socio-demographic variables were included as independent variables. These variables include age, sex, educational status, place of residence, patient's delay and types of TB.

Results

A total of 384 TB patients participated in this study – 225(58.6%) males and 159(41.4%) females. Table 1 explains the four dimensions regarding the knowledge of TB. The majority of patients had knowledge about the symptoms of TB. The most frequently reported knowledge of symptoms was related to coughing (97.9%), followed by chest pain (60.2%), shortness of breath (24.2%), loss of appetite (50.8%), weight loss (56.8%) and fever with night sweating (70.6%). Most of the respondents were aware of the transmission routes of TB. Almost all

(94.0%) respondents knew TB is transmitted through sneezing and 89.8% respondents were aware that TB is communicable. Most of the patients were found to have good knowledge about treatment of TB. Essentially all of the respondents (99.0%) were knowledgeable that treatment is available, 95.1% of respondents understood treatment is free and 97.1% believed regular intake of medicine can cure them. Close to half of the respondents (48.4%) claimed that irregular intake of medicine can cause death. Almost all of the respondents (98.4%) knew staying far away from TB affected people was one of the fundamental preventive measures. Almost one third of respondents (30.7%) had unsatisfactory knowledge about the vaccination for TB.

Table 1. Distribution of Knowledge Related Variables and Their Categories

Dimensions	Response variables	Categories	Frequency (%)
Dimension of Knowledge about Symptoms (DKS)	Coughing with and without blood more than 3 weeks	0= No 1=Yes	8 (2.1%) 376 (97.9%)
	Chest pain	0= No 1=Yes	153 (39.8%) 231 (60.2%)
	Shortness of breath	0= No 1=Yes	291 (75.8%) 93 (24.2%)
	Loss of appetite	0= No 1=Yes	189 (49.2%) 195 (50.8%)
	Weight loss	0= No 1=Yes	166 (43.2%) 218 (56.8%)
	Fever with night sweating	0= No 1=Yes	113 (29.4%) 271 (70.6%)
	Is TB communicable	0= No 1=Yes	39 (10.2%) 345 (89.8%)
	Through sneezing	0= No 1=Yes	23 (6.0%) 361 (94.0%)
Dimension of Knowledge about Transmission (DKT)	Through air	0= No 1=Yes	129 (33.6%) 255 (66.4%)
	Through using things of affected people	0= No 1=Yes	203 (52.9%) 181 (47.1%)
	Is treatment available	0= No 1=Yes	4 (1.0%) 380 (99.0%)
	Is treatment free	0= No 1=Yes	19 (4.9%) 365 (95.1%)
Dimension of Knowledge about Treatment (DKTr)	Is regular intake of medicine cure patient	0= No 1=Yes	11 (2.9%) 373 (97.1%)
	Is irregular intake of medicine cause death/MDR	0= No 1=Yes	198 (51.6%) 186 (48.4%)
	Is vaccine available	0= No 1=Yes	266 (69.3%) 118 (30.7%)
	Stay far away from affected people when sneeze	0= No 1=Yes	20 (5.2%) 364 (94.8%)
Dimension of Knowledge about Prevention (DKP)	Not use things of affected people	0= No 1=Yes	172 (44.8%) 212 (55.2%)
	Get always clean	0= No 1=Yes	68 (17.7%) 316 (82.3%)
	Cronbach's Alpha	0.785	

The mean knowledge levels (MKL) of TB patients are presented in Table 2. The pulmonary TB patients had a better knowledge level than the extra-pulmonary TB patients. The mean OKI for pulmonary patients was 0.762 and for extra-pulmonary patients was 0.572. On the basis of patient's type, the patients recorded as old cases had higher MKL for all of the dimensions as compared to the new cases. The mean OKI for old cases was 0.763 and 0.701 of for new cases.

Table 2. Distribution of Mean Knowledge by TB Types and Patient Types

Dimensions	Types and categories of patients		Mean
Dimension of Knowledge about Symptoms (DKS)	Patient's type	Extra-Pulmonary	0.363
		Pulmonary	0.684
	Patient's Category	New case	0.577
		Old case	0.704
Dimension of Knowledge about Transmission (DKT)	Patient's type	Extra-Pulmonary	0.575
		Pulmonary	0.802
	Patient's Category	New case	0.731
		Old case	0.795
Dimension of Knowledge about Treatment (DKTr)	Patient's type	Extra-Pulmonary	0.777
		Pulmonary	0.874
	Patient's Category	New case	0.837
		Old case	0.901
Dimension of Knowledge about Prevention (DKP)	Patient's type	Extra-Pulmonary	0.575
		Pulmonary	0.686
	Patient's Category	New case	0.658
		Old case	0.651
Overall Knowledge Index (OKI)	Patient's type	Extra-Pulmonary	0.572
		Pulmonary	0.762
	Patient's Category	New case	0.701
		Old case	0.763

Table 3 illustrates patients' socio-demographic characteristics affecting knowledge level. The study results revealed that more than half (61.8%) of all male respondents, and half (50.3%) of female respondents had sufficient knowledge about TB. The MKL was lower among females (0.684) than of males (0.732) and was higher in ages 21-35 years old (0.774) as compared to other age groups. Moreover, respondents in the 21-35 year age group had the highest percentage with sufficient knowledge (71.0%). Around half of the respondents (52.6%) were less educated, and the percentage of respondents with sufficient knowledge was highest among the most educated group, those with 12 or more years of schooling (68.5%). Most of the patients (61.5%) lived in the urban area. More than three-fourth patients (76.3%) were delayed in diagnosis of their TB status. Pulmonary patients were found to have a higher percentage (74.0%) among all patients, and most of them (70.8%) had had sufficient knowledge. The χ^2 test results suggest that sex, age, educational status, and TB type are significantly associated with the knowledge level.

Table 3. Distribution of Knowledge Level According to Different Socio-Demographic Variables and Associations of Knowledge Level with Such Variables

Explanatory variables	Mean knowledge level	Knowledge level		Total
		Moderate knowledge	Sufficient knowledge	
Sex**				
Male	0.732	86 (38.2%)	139 (61.8%)	225 (58.6%)
Female	0.684	79 (49.7%)	80 (50.3%)	159 (41.4%)
Age***				
< 20 years	0.674	25 (59.5%)	17 (40.5%)	42 (10.9%)
21-35 years	0.774	40 (29.0%)	98 (71.0%)	138 (35.9%)
36-50 years	0.702	57 (46.7%)	65 (53.3%)	122 (31.8%)
> 51 years	0.643	43 (52.4%)	39 (47.6%)	82 (21.4%)
Educational status**				
0-5 years of schooling	0.666	101 (50.0%)	101 (50.0%)	202 (52.6%)
6-12 years of schooling	0.760	47 (36.7%)	81 (63.3%)	128 (33.3%)
> 12 years of schooling	0.770	17 (31.5%)	37 (68.5%)	54 (14.1%)
Residence				
Rural	0.714	59 (39.9%)	89 (60.1%)	148 (38.5%)
Urban	0.711	106 (44.9%)	130 (55.1%)	236 (61.5%)
Patient's delay				
No	0.696	42 (46.2%)	49 (53.8%)	91 (23.7%)
Yes	0.717	123 (42.0%)	170 (58.0%)	293 (76.3%)
TB types***				
Extra-Pulmonary	0.572	82 (82.0%)	18 (18.0%)	100 (26.0%)
Pulmonary	0.762	83 (29.2%)	201 (70.8%)	284 (74.0%)
	0.712	165 (43.0%)	219 (57.0%)	384 (100.0%)

Note: *** $p < 0.01$; ** $p < 0.05$.

Table 4 presents the results of the multivariate analysis of the factors affecting knowledge level of the TB patients. The results of the logistic regression suggest that the socio-demographic variables selected for the analysis are generally important predictors of knowledge level of the TB patients. Female TB patients were 0.824 times less likely to be informed compared to their male counterpart. Overall, age has a significant positive effect on knowledge level. The patients aged 21-35 years were 3.660 times more likely to have sufficient knowledge of TB than those who are 20 years of age or younger. Large, statistically significant differences in knowledge level among TB patients were observed by educational level. Respondents with the highest level of education (12 years or more of schooling) were 8.097 times more likely to have sufficient knowledge compared to those who completed 0-5 years of schooling. Urban respondents were 1.123 times more aware of TB compared to rural respondents. The patients with delayed diagnosis were 0.89 time less likely to be informed. In case of TB type of the patients, pulmonary TB patients were 26.827 times more likely to be informed than extra-pulmonary TB patients.

Table 4. Affecting Factors on Knowledge Level of the TB Patients

Explanatory variables	Beta (β) values	S.E of β	Odd ratios (OR)	95% CI	
				Upper level	Lower level
Sex					
Male (RC)	-0.194	0.273	1.000 0.824	0.482	1.407
Age					
<20 years (RC)			1.000		
21-35 years	1.297***	0.471	3.660	1.453	9.216
36-50 years	0.570	0.487	1.769	0.681	4.598
>51 years	0.410	0.512	1.507	0.553	4.109
Educational status					
0-5 years of schooling (RC)			1.000		
6-12 years of schooling	1.426***	0.366	4.163	2.032	8.527
>12 years of schooling	2.091***	0.519	8.097	2.929	22.382
Residence					
Rural(RC)			1.000		
Urban	0.116	0.288	1.123	0.638	1.976
Patient's delay					
No (RC)			1.000		
Yes	-0.117	0.323	0.890	0.472	1.675
TB types					
Extra-Pulmonary (RC)			1.000		
Pulmonary	3.289***	0.401	26.827	12.216	58.915
Constant	-3.497	.705	.030		
-2 × Log likelihood	384.045				
Cox & Snell R^2	0.307				

Note: *** $p < 0.01$; CI=Confidence Interval; RC=Reference Category; S.E=Standard Error.

Discussion

The objective of this study was to determine the knowledge level of the TB patients and to what extent socio-demographic factors contribute to differences in knowledge level in Rajshahi city, Bangladesh. We collected and analyzed data from 384 TB patients who are diagnosed with TB and receiving anti-TB treatment. To our knowledge, no previous study has examined the knowledge level in Rajshahi city, Bangladesh. The findings of this study suggest that more than 50% of the respondents knew about the symptoms of TB. Our findings suggest that socio-demographic variables played important roles on the knowledge level of the TB patients.

Logistic regression analysis showed age, educational status, and patient type were significantly associated with TB knowledge level. Patients in the 21-35 years of age group were significantly more likely to have sufficient knowledge about TB ($p < 0.01$) than those of < 20 years of age group. Likewise, the patients who completed 6-12 years of schooling and > 12 years of schooling were also significantly more likely to have sufficient knowledge status ($p < 0.01$) than those who completed 0-5 years of schooling. Hoa et al. (2004) found similar results, which, consistent with our findings, highlighted educational level as an important determinant of people's level of knowledge of TB, and persons with a higher level of education scored better on TB knowledge indices than those with less education or those who were illiterate. This study has shown that, while socio-demographic factors may not necessarily be causative factors to becoming infected with TB, they do significantly influence knowledge levels of TB patients.

The findings of this study are consistent with some other national studies. For example, one study conducted in Iraq found the similar result with around 50% of the patients had good knowledge about the symptoms of TB (Wang et al. 2008). Another study conducted in Ndola,

Zambia found that over 70% of TB patients had an idea about the symptoms, including fever with night sweating, weight loss, loss of appetite and coughing with and without blood for more than 3 weeks (Hashim et al. 2003). Coughing with and without blood for more than 3 weeks is regarded as a TB suspicious symptom and only 16% people in Yangzhong County, a rural area of China, had heard about it (Kaona et al. 2004). However, in this study 97.9% patients knew coughing with and without blood for more than 3 weeks is one of the major symptoms of TB. Another study conducted in the north and central regions of Vietnam found coughing was the most frequently mentioned symptom, and was reported by 298 patients (81.9%) (Hoa et al. 2004). A similar result was found in Ibadan, Nigeria where a study conducted among pulmonary patients revealed more than 80% of the patients recognized coughing as major symptoms of TB (Fatiregun and Ejekam 2010). Focused health education programs in Bangladesh appear to have made a great impact on the level of patient knowledge about TB.

Since TB is a contagious, communicable disease; it spreads through contact with an infected person, making the understanding of human transmission of infection absolutely critical to its control (Allotey and Gyapong 2008). Around 95.3% of pulmonary patients in the north and central regions of Vietnam were aware that TB is a contagious disease (Mondal et al. 2014). From the current study, we find almost all patients have heard about TB and more than 89.8% knew it was a communicable disease, and almost all patients knew the treatment of TB is available, free, and regular intake of medicine can cure a patient. Nearly half of the patients had an idea that contact with infected people's belongings can spread TB. Our findings are consistent with several other studies which have investigated knowledge levels of TB in other countries (Wang et al. 2008; Kaona et al. 2004; Hoa et al. 2004; Fatiregun and Ejekam 2010).

While overall knowledge levels are relatively high in Bangladesh, specific knowledge about TB associated with health-care seeking behaviors still appears unsatisfactory. Many people living with TB remain underreported because of lack of knowledge about treatment and the TB control program. Poor knowledge of TB patients concerning their disease may contribute to the high prevalence of TB disease in the country (Khan et al. 2006). Our findings suggest that the mean level of knowledge was higher among pulmonary patients because most of the symptoms, transmission, and prevention items listed in this study were related to the pulmonary patients. This fact highlighted the difference in knowledge level among pulmonary TB patients and other TB patients. The study also showed the mean knowledge level among the patients identified as new cases was lower than the patients recorded old cases. Specifically, the mean OKI for new cases was 0.701 and for old cases was 0.763. This result suggests that overall knowledge levels are relatively high for patients of both categories, and are similar to the Vietnam study, which showed patients in the north and middle of Vietnam had reasonably sufficient knowledge about TB (Fatiregun and Ejekam 2010). However, several other studies have concluded that knowledge about TB among new pulmonary TB patients was quite low, which is contradicted in this study (Liam et al. 1999; Bhat et al. 1999).

A prior study conducted in Hyderabad, India found socio-demographic variables, which included age, sex, educational status, occupation, monthly income, nutritional status, addiction etc., are closely linked with TB prevalence and knowledge level (Fatiregun and Ejekam 2010; Soomro and Qazi 2009; Ahsan et al. 2004). In the present study, we found that age, sex, and educational status were variables that influence TB related knowledge. This result was similar to the study conducted among patients in Iraq, which highlighted age and educational status as being significant predictors of knowledge of TB (Kaona et al. 2004). In our study, a large number of patients seeking treatment at the study sites were males, of which knowledge level was high (0.732). A number of other studies have shown that, in developing countries, male TB patients account for two thirds of total reported TB patients (Nakagawa et al. 2001). Globally, approximately 70% more males are notified of smear-positive TB tests than females

(Diwan and Thorson 1999). According to WHO, the prevalence of TB is more common among men than women (Neyrolles and Quintana 2009). Ahsan et al. (2004) found that female patients are mostly illiterate and live in poor socio-economic conditions. It is quite apparent that socio-economic and cultural factors are important determinants of gender differentials in TB prevalence (Nakagawa et al. 2001). Females are often unable to reach health facilities because a woman's position in the household, economic dependence, and illiteracy would be restricting factors (Kaulagekar and Radkar 2007).

Aside from gender differences, our study also revealed that most of the TB patients seeking treatment belonged to the 21-35 years of age group, and that the MKL (0.774) was higher for this age group. Additionally, higher MKL was found among the patients completing >12 years of schooling. These results suggest that age and education level are important determinants of general knowledge of TB, and this finding is also consistent with other studies (Fatiregun and Ejekam 2010; Bhat et al. 1999). It should also be noted, however, that this study has some limitations. It is based on a cross sectional data and sample size is small. The population consists of Rajshahi city and not of Bangladesh as an entire country. Given the limitations of the present study, future studies should consider larger sample, especially, nationally representative sample.

Conclusion

Knowledge level among the TB patients in Rajshahi city is relatively high, yet it was dependent on some socio-demographic factors. Males were better informed than females, and the young adult population, ages 21-35, was more aware about TB transmission routes and preventative measures. Also, more highly educated and urban area patients were comparatively well informed about TB infection. Patients with greater knowledge about TB were also less likely to be delayed in seeking diagnosis of their TB status. Strengthening awareness of TB and improving the accessibility of healthcare services is essential in TB control strategies, especially under the current vertical TB control system.

Ethical considerations

Ethical issues (Including plagiarism, Informed Consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc.) have been completely observed by the authors.

Acknowledgements

The authors are very grateful to the Department of Population Science and Human Resource Development, University of Rajshahi, Bangladesh by giving an opportunity to complete this study fruitfully. Authors are very grateful to the respondents as well as the authority of the TB centers. Thanks are also due to the editor and reviewers of the Journal of African Health Sciences for their valuable comments and criticisms, which led to a greatly improved revision of this paper.

References

- Ahsan, G., Ahmed, J., Singhasivanon, P., Kaewkungwal, J., Okanurak, K., Suwannapong, N., Akarasewi, P., Majid, M. A., Begum, V., and Belayetali, K. 2004. "Gender Difference in Treatment Seeking Behaviors of Tuberculosis Cases in Rural Communities of Bangladesh." *Southeast Asian Journal of Tropical Medicine and Public Health* 35(1): 126-135.
- Allotey P., and Gyapong M. 2008. "Gender in Tuberculosis Research." *International Journal of Tuberculosis and Lung Disease* 12(7): 831-836.
- Bhat, S., Singal, N., Aggarwal, C. S., and Jain R. C. 1999. "Knowledge, Attitudes and Practices of Newly Diagnosis Sputum Positive Cases of Pulmonary Tuberculosis." *Journal of Communicable Diseases* 31(4): 247-252.
- Demissie, M., Lindtjorn, B., and Berhane, Y. 2002. "Patient and Health Service Delay in the Diagnosis of Pulmonary Tuberculosis in Ethiopia." *BMC Public Health* 2: 23.
- Diwan, V. K., and Thorson, A. 1999. "Sex, Gender, and Tuberculosis." *Lancet* 353(9157): 1000-1001.
- Fatiregun, A. A., and Ejekam, C. C. 2010. "Determinants of Patients Delay in Seeking Treatment among Pulmonary Tuberculosis Cases in a Government Specialist Hospital in Ibadan, Nigeria." *Tanzania Journal of Health Research* 12(2): 1-9.
- Gupta, R. K., Gupta, A., Jamwal, D. S., and Suri, S. P. 2002. "A Socio-Epidemiological Study of Tuberculosis in a Rural Area." *JK Science: Journal of Medical Education & Research* 4(3): 119-122.
- Hashim, D. S., Kubaisy, W. A., and Dulayme, A. A. 2003. "Knowledge, Attitudes and Practices Survey among Health Care Workers and Tuberculosis Patients in Iraq." *Eastern Mediterranean Health Journal* 9(4): 718-731.
- Hoa, N. P., Diwan, V. K., Co, N. V., and Thorson, A. E. K. 2004. "Thorson AEK: Knowledge about Tuberculosis and its Treatment among New Pulmonary TB Patients in the North and Central Regions of Vietnam." *International Journal of Tuberculosis and Lung Disease* 8(5): 603-608.
- Karim, F., Islam, M. A., and Chowdhury, A. M. 2007. "Gender Differences in Delays in Diagnosis and Treatment of Tuberculosis." *Health Policy and Planning* 22(5): 329-334.
- Kaona, F. A. D., Tuba, M., and Sisiya, S., and Sikaona, L. 2004. "An Assessment of Factors Contributing to Treatment Adherence and Knowledge of TB Transmission among Patients on TB Treatment." *BMC Public Health* 4: 68
- Kaulagekar, A., and Radkar, A. 2007. "Social Status Makes a Difference: Tuberculosis Scenario during National Family Health Survey-2." *Indian Journal of Tuberculosis* 54(1): 17-23.

- Khan, J. A., Irfan, M., and Zaki A, Beg, M., Hussain, S. F., and Rizvi, N. 2006. "Knowledge, Attitude and Misconceptions regarding Tuberculosis in Pakistani Patients." *Journal of Pakistan Medical Association* 56(5): 211-214.
- Koay, T. K. 2004. "Knowledge and Attitudes towards Tuberculosis among the People Living in Kudat District, Sabah." *Medical Journal of Malaysia* 59(4): 502-511.
- Liam, C. K., Lim, K. H., Wong, C. M. M., and Tang, B. G. 1999. "Attitudes and Knowledge of Newly Diagnosed Tuberculosis Patients regarding the Disease, and Factors Affecting Treatment Compliance." *International Journal of Tuberculosis and Lung Disease* 3(4): 300-309.
- Long, Q., Li, Y., and Wang, Y. 2008. "Barriers to Accessing TB Diagnosis for Rural-to-Urban Migrants with Chronic Cough in Chongqing, China: A Mixed Methods Study." *BMC Health Services Research* 8: 202.
- Mauch, V., Woods, N., Kirubi, B., Kipruto, H., Sitienei, J., and Klinkenberg, E. 2011. "Assessing Access Barriers to Tuberculosis Care with the Tool to Estimate Patients' Costs: Pilot Results from two Districts in Kenya." *BMC Public Health* 11: 43.
- Mondal, M. N. I., and Shitan, M. 2013a. "Impact of Socio-Health Factors on Life Expectancy in the Low and Lower Middle Income Countries." *Iranian Journal of Public Health* 42(12): 1354-1362.
- Mondal, M. N. I., and Shitan, M. 2013b. "Factors Affecting the HIV/AIDS Epidemic: An Ecological Analysis of Global Data." *African Health Sciences* 13(2): 294-303.
- Mondal, M. N. I., Chowdhury, R., and Sayem, A. 2014. "Associated Factors of Pulmonary Tuberculosis in Rajshahi City of Bangladesh." *Journal of Human Ecology* 45(1): 61-68.
- Nakagawa, M. Y., Ozasa, K., Yamada, N., Osuga, K., Shimouchi, A., Ishikawa, N., Bam, D. S., and Mori, T. 2001. "Gender Difference in Delays to Diagnosis and Health Care Seeking Behavior in a Rural Area of Nepal." *International Journal of Tuberculosis and Lung Disease* 5(1): 24-31.
- Neyrolles, O., and Quintana, M. L. 2009. "Sexual Inequality in Tuberculosis." *PLOS Medicine* 6(12): e1000199.
- Ngadaya, E. S., Mfinanga, G. S., Wandwalo, E, and Morkve, O. 2009. "Delay in Tuberculosis Case Detection in Pwani region, Tanzania: A Cross-sectional Study." *BMC Health Services Research* 9: 196.
- Soomro, J. A., and Qazi, H. A. 2009. "Factors Associated with Relapsed Tuberculosis in Males and Females: A Comparative Study." *Tanaffos* 8(3): 22-27.
- United Nations Development Program (UNDP). 2011. "Human development Report 2011." http://hdr.undp.org/sites/default/files/reports/271/hdr_2011_en_complete.pdf.

Wang, J., Fei, Y., Shen, H., and Xu, B. 2008. "Gender Difference in Knowledge of Tuberculosis and Associated Health-care Seeking Behaviors: A Cross-sectional Study in a Rural Area of China." *BMC Public Health* 8: 354. .

World Health Organization (WHO). 2011. "Global tuberculosis control: WHO report 2011." http://apps.who.int/iris/bitstream/10665/44728/1/9789241564380_eng.pdf.

Zafar Ullah, A. N., Newell, J. N., Ahmed, J. U., Hyder, M. K. A., and Islam, A. 2006. "Government-NGO Collaboration: the Case of Tuberculosis Control in Bangladesh." *Health Policy and Planning* 21(2): 143-155.

Author Biography

Nazrul Hoque (Ph.D., Pennsylvania State University) is a Senior Researcher at the Hobby Center for Public Policy at the University of Houston, Texas. He is an applied demographer whose research interests are in the areas of population estimates and projections, fertility, mortality, migration, and aging. He is particularly interested in examining the impact of future demographic changes on different types of public services and policy issues. He has produced official annual population estimates for the State of Texas and its 254 counties and nearly 2,000 places and biennial population projections for the State of Texas and 254 Counties in Texas by age, sex, and race/ethnicity from 1989 to 2013. He is a Fulbright scholar. His research has been published in the journals of *Population Research and Policy Review*, *Canadian Studies in Population*, *Canadian Journal on Aging*, *Spatial Economic Analysis*, *World Journal of Aids*, *Demography India*, and *African Health Sciences*. He has co-authored and co-edited books in his areas of expertise.